

Modelling and Forecasting of Dissolved Oxygen Using the ARIMA Model: The case of the Ledrat Dam (Medea Province- Algeria)

Mostefaoui Houda ¹, Messaoudi Ayoub ²

¹ Laboratory of Biotechnology and Valorization of Biological Resources. Department of Agronomy. Faculty of Sciences. University of Medea. Ouzra 26100, Medea 26100, Algeria. Mail: mostefaoui.houda@univ-medea.dz

² Laboratory of Biotechnology and Valorization of Biological Resources. Department of Agronomy. Faculty of Sciences. University of Medea. Ouzra 26100, Medea 26100, Algeria. Mail: messaoudi.ayoub@univ-medea.dz

Received: 10-12-2025

Accepted: 25-02-2026

Published: 126-04-2026

SUMMARY:

Dissolved oxygen (DO) is a key indicator of water quality. This study focuses on the modelling of the temporal evolution of dissolved oxygen (DO) mg/L in the waters of the Ledrat dam, located in the wilaya of Medea, using the ARIMA model. The descriptive analysis shows significant variability in the dissolved oxygen values during the period of 2012-2022 with seasonal fluctuations and outliers greater than 12 mg/L, as well as the decomposed time series shows a nonlinear trend. The ARIMA model shows a good performance with an RMSE of 1.4191, an MAE of 1.0727 and a MAPE of 13.28%, which indicates adequate accuracy for the forecasts. The model validation using the Ljung-Box test ($Q^*=22.448$, $p\text{-value}=0.2625$) confirms the independence of the residues, attesting to the quality of the fitted model. The results obtained prove that the ARIMA model is effective in capturing dissolved oxygen (DO) levels and providing relevant forecasts that can be used as a decision support tool to support the optimized management of the dam's water resources and preserve water quality.

KEYWORDS: dissolved oxygen, ARIMA model, Ljung-Box test, Ledrat Dam.

INTRODUCTION:

Water, a vital commodity on Earth, is a recyclable resource. It is essential for the development of all areas of human effort (Obi et al., 2006; Merrah, 2010; Messaoudi et al., 2026)

Rapid urbanization, industrial expansion and intensification of agricultural practices have significantly increased the pollution of aquatic environments, both chronic and sporadic. Many rivers and reservoirs are now experiencing a continuous deterioration in their physico-chemical quality, under the effect of sustained anthropogenic pressure linked to domestic, industrial and agricultural discharges (Benoit and Comeau, 2005; Bekkoussa et al., 2013; Abidi et al., 2014; Benabdelmalek et al., 2020).

Among the indicators of ecological health, dissolved oxygen (DO) is a fundamental parameter. It reflects the metabolic balance of aquatic ecosystems, resulting from the trade-off between photosynthetic production and the respiratory demand of biomass (Wetzel, 2001; Boyd, 2020). Maintaining sufficient oxygen levels is essential for the respiration of most organisms (fish, macroinvertebrates, and aerobic bacteria).

When DO concentrations fall below the critical threshold of 2mg/L, the waters are classified as hypoxic or even anoxic. This phenomenon represents a severe threat to biodiversity and leads to a profound alteration of biogeochemical cycles. The absence of oxygen promotes the release of nutrients and toxic compounds such as H_2S , Fe^{2+} , and Mn^{2+} from sediments (Wetzel, 2001; Rabalais et al., 2010; Jane et al., 2021; Li et al., 2023). These processes aggravate internal eutrophication and drastically reduce the biological richness of the environment (Nürnberg, 2002; Beutel, 2003; Søndergaard et al., 2017).

Episodes of hypoxia are particularly common in deep dams. During the summer, thermal stratification isolates the deep layers (hypolimnion), preventing their reoxygenation and promoting an increased benthic consumption of available oxygen (Kraemer et al., 2021; Zhang et al., 2022). This phenomenon drastically limits the oxygen renewal of bottom water (Amieva et al., 2023). Faced with the complexity of DO fluctuations influenced by nycthemeral cycles, terrigenous organic inputs and climate changes that accelerate deoxygenation (Kraemer et al., 2021).

Modelling appears to be a crucial decision-making tool. The application of statistical time-series models, such as the Autoregressive Integrated Moving Average (ARIMA) models, offers significant potential for predicting

changes in water quality. These tools support ecological management in accordance with the requirements of the Water Framework Directive, which recommends thresholds of 5-6 mg/L to ensure good environmental status (European Commission, 2018). In this perspective, the present work aims to analyze a time series of dissolved oxygen over a period of 10 years at the Ledrat dam.

This study aims to analyze the time series of dissolved oxygen in order to understand its dynamics, and to identify seasonal variations and then to develop an ARIMA model allowing the modeling and forecasting of this essential parameter of water quality.

MATERIALS AND METHODS:

Study site

The Ledrat dam is located in the commune of Omaria, wilaya of Medea (fig.01), in the north of Algeria (35°58'N; 2°48'E). Commissioned in 1987, it is one of the major hydraulic structures in the catchment area of the Chiffa wadi, a tributary of the Medjerda river basin. With an initial storage capacity estimated at 85 million m³, this structure plays a strategic role in the supply of water for irrigation, local industry and human consumption in several municipalities in the wilaya of Médéa (ANRH, 2020; Benabdelmalek *et al.*, 2020).




Figure 01: Map of the geographical location of the Ledrat Dam

Sampling Method:

Monthly measurements of dissolved oxygen (DO) are collected over a period of 10 years by the National Agency for Water Resources (ANRH, Station de Médéa.)

Water samples were collected monthly over a 10-year period (2012–2022)

Analytical methods:

Measurements of physicochemical parameters were performed *in situ* according to the standardized protocols of the APHA (American Public Health Association, 2017) and Rodier *et al.* (2021). Dissolved oxygen measurements were performed *in situ* using a portable HACH HQD DR-1900 **multiparameter probe** equipped with an optical LDO (Luminescent Dissolved Oxygen) probe. This optical sensor has the advantage of high accuracy (± 0.1 mg/L or $\pm 1\%$ of the measured value), low drift and no oxygen consumption during measurement, unlike conventional electrochemical probes (HACH, 2020; Rode *et al.*, 2016; Li *et al.*, 2023; Boyd, 2020).

Statistical analysis

The data obtained were statistically processed to identify trends and monthly variations in the physicochemical parameters of dissolved oxygen (DO). The data analysis was carried out using the R software (CRAN, 2024).

RESULTS AND DISCUSSIONS

The results of the analysis of the behaviour of the dissolved oxygen through two complementary approaches (fig.02), show that the dissolved oxygen values are generally centered around 8mg/L, with moderate variability and some outlier values above 12 mg/L.

The decomposition of the time series by the STL method highlights a general upward trend until 2018, followed by a marked and continuous decline between 2019 and 2022. This trend reversal suggests a recent deterioration in the dam's water quality, potentially related to increased organic loading, accelerated sedimentation, or the effects of climate change on water stratification and temperature (Søndergaard *et al.*, 2017; Zhang *et al.*, 2022; Li *et al.*, 2023; Jane *et al.*, 2021 ; Kraemer *et al.*, 2021).




Figure 02: Descriptive analysis and decomposition of the oxygen time series below

The seasonal component is particularly pronounced, confirming the dominant influence of annual cycles on dissolved oxygen dynamics. The highest concentrations are generally observed in winter and spring, while minima are recorded in late summer and autumn, a period of thermal stratification and high bacterial decomposition activity (Wetzel, 2001; Nürnberg, 2002; Jane *et al.*, 2021; Box *et al.*, 1994 ; Yue *et al.*, 2002; Messaoudi *et al.*, 2026).

Finally, the analysis of decomposition residues shows relatively low and random values, indicating that the majority of the observed variations in DO concentrations are explained by the interannual trend and the seasonal cycle (Rabalais *et al.*, 2010 ; Søndergaard *et al.*, 2017 ; Li *et al.*, 2023). These results highlight the need for enhanced monitoring of the recent decline in dissolved oxygen, as prolonged hypoxia could lead to serious ecological consequences, such as fish mortality, reduced aquatic biodiversity, and leaching of nutrients and heavy metals from sediments (Rabalais *et al.*, 2010 ; Søndergaard *et al.*, 2017; Kraemer *et al.*, 2021).

ARIMA MODEL RESULTS:

Verification of the Stationarity of the Data:

Before applying the ARIMA model, it is imperative to verify the stationarity of the time series, which is a fundamental condition for ensuring the statistical validity and predictive reliability of such models (Box *et al.*, 2015; Hyndman and Athanasopoulos, 2018; Box *et al.*, 1994; Shumway & Stoffer, 2017). Indeed, nonstationarity can lead to biased estimates, unreliable forecasts, and misinterpretation of temporal relationships (Shumway and Stoffer, 2017; Hyndman & Athanasopoulos, 2018).

For this purpose, the augmented Dickey-Fuller test (ADF) was used. This test, widely used in hydrology and limnology for the analysis of water quality time series (Hirsch & Slack, 1984; Yue *et al.*, 2002; Zhang *et al.*,

2022), allows to test the null hypothesis (H_0) according to which the series has a unit root (non-stationary process) against the alternative hypothesis (H_1) of stationarity (Dickey and Fuller, 1979; Said and Dickey, 1984). The results obtained (p -value = 0.022 < 0.05) lead to the rejection of the null hypothesis at the 5% significance level. This statistic leads to the conclusion that the dissolved oxygen time series is stationary (at least in mean and variance). This statistical property ensures that the parameters of the ARIMA model (orders p , d , q) can be estimated robustly and that these coefficients will remain stable over time, a guarantee of reliability for short-term forecasts (Box et al., 1994; Parviz et al., 2010; Enders, 2015; Cryer and Chan 2008).

Model Results:

One of the most important and widely used time series models is the Box-Jenkins modeling approach, commonly referred to as the Autoregressive Integrated Moving Average (ARIMA) (Box et al., 1994).

ARIMA models are flexible in that they can represent several different time series, that is to say, pure autoregressive (AR), pure moving average (MA), and combination AR and MA (ARMA) series, although they are limited by their assumed linear form (Parviz et al., 2010).

Table 01 : ARIMA model results

Metric	Value
Sigma^2	2.11
AIC	480.14
AICc	480.81
BIC	497.39
ME	-0.0516
RMSE	1.4191
MFA	1.0727
MPE (%)	-3.3944
MAPE (%)	13.2822
MASE	0.7073
ACF1	-0.0082

Note: Sigma^2 : residual variance; **AIC**: akaike information criterion; **AICc**: Corrected Akaike Information Criterion; **BIC**: Bayesian information criterion; **ME**: mean error; **RMSE**: root mean squared error; **MAE**: mean absolute error; **MPE**: mean percentage error; **MAPE**: mean absolute percentage error; **MASE**: mean absolute scaled error; **ACF1**: autocorrelation function at lag 1

Analysis of the results obtained for the ARIMA model (3,1,1) (1,0,0), applied to dissolved oxygen (DO) data, indicates a robust and well-fitting model. Fit metrics, such as AIC (480.14), AICc (480.81), and BIC (497.39), show reasonably low values, indicating that the model achieves a good trade-off between complexity and accuracy. The prediction errors (MOE = -0.0516, RMSE = 1.4191, MAE = 1.0727) are small, which reinforces the idea that the model effectively captures the variations of the series. The p -value of 0.2625 from the Ljung-Box test confirms that the model residuals do not have significant autocorrelation, which is essential for a well-fitting ARIMA model.

Residue Analysis and Model Validation:

The validation of the ARIMA model (3,1,1) (1,0,0) [12] was carried out through several statistical and graphical diagnoses in order to verify the quality of the fit and the absence of significant residual structure.

Table 02 : Model Validation

Metric	Value
Q^*	22.448
Degrees of Freedom (df)	19.0
P-value	0.2625
Model Degrees of Freedom	5.0
Total Delays Used	24.0

The Ljung-Box test was applied to the model residues. The results (Tab. 2) show a Q^* value of 22.448 for 19 degrees of freedom, associated with a p-value of 0.2625. Since this value is above the significance level of 0.05, the null hypothesis of the absence of significant autocorrelation of the residues cannot be rejected (Ljung and Box, 1978; Box *et al.*, 2015; Hyndman & Athanasopoulos, 2018). This result indicates that the residues behave like white noise, an essential condition for validating an ARIMA model (Hyndman and Athanasopoulos, 2018; Shumway and Stoffer 2017).




Figure 03: Analysis of model residuals

Graphical examination of the residues (Figure 4) shows random fluctuations around zero with no systematic trend or residual cyclic structure, in accordance with the visual validation criteria recommended for environmental time series (Hirsch & Slack, 1984; Yue *et al.*, 2002; Cryer and Chan, 2008). The analysis of the autocorrelation function (ACF) of the residues confirms that all autocorrelation coefficients remain included in the 95% confidence intervals, showing the absence of significant serial correlation (Brockwell & Davis, 2016). On the other hand, the examination of the distribution of the residues (histogram and Quantile-Quantile diagram) indicates a reasonable approximation to a normal distribution, although slight asymmetries are observable. This near-normality of the residues is important because it ensures the validity of the confidence intervals and statistical tests associated with the model (Shumway and Stoffer, 2017; Brockwell and Davis, 2016).

In summary, all the statistical and graphical diagnoses confirm that the ARIMA model (3,1,1)(1,0,0) is adequate for modelling and predicting dissolved oxygen levels in the Ledrat dam, particularly in the context of dissolved oxygen monitoring and anticipation of hypoxia risks (Amieva *et al.*, 2023; Zhang *et al.*, 2022).

Forecasts:

Figure (04) illustrates the predictions of the dissolved oxygen values by the ARIMA model.

It can be seen that the observed values show significant variability over the period studied, while the forecasts indicate a relatively stable evolution around 7-8 mg/L.

The forecast intervals indicate that there is significant uncertainty, related to the natural variability of dissolved oxygen and the fluctuations observed in the time series.




Figure 04: Predictions of dissolved oxygen values (DO2)

CONCLUSION:

In conclusion, the ARIMA model (3,1,1) (1,0,0) [12] has proven to be adequate and robust for modelling and predicting the spatio-temporal variations of dissolved oxygen (DO) in the waters of the Ledrat dam over the period 2012–2022. The fit criteria (AIC, AICc, BIC), the residue analysis, the Ljung-Box test ($p = 0.2625$) as well as the examination of the autocorrelation functions confirm that the model effectively captures the temporal structure of the series, both in its trend and seasonal components, without significant residual autocorrelation or systematic bias.

The good performance of the model, illustrated by moderate prediction errors (RMSE = 1.42 mg/L; MAE = 1.07 mg/L; MAPE = 13.28%), demonstrates its reliability in making short- and medium-term forecasts. These forecasts are a valuable operational tool for managers, allowing them to anticipate hypoxia episodes, identify critical periods and implement appropriate preventive measures.

However, the marked decline in dissolved oxygen observed between 2019 and 2022 raises concerns about the gradual deterioration of the dam's water quality. This phenomenon, likely amplified by climate change, thermal stratification and anthropogenic inputs, could lead to important ecological consequences, such as fish mortality, nutrient release from sediments and the development of cyanobacteria.

Ultimately, this work demonstrates the interest of ARIMA modelling approaches combined with long time series analysis for the monitoring and sustainable management of reservoirs. It paves the way for the integration of these models with remote sensing and machine learning approaches, to improve the prediction and proactive management of water quality in Algerian dams.

BIBLIOGRAPHIC REFERENCE:

- Abidi, Selma, Boualem Remini and Lakhdar Kini. 2014. "Water quality of the main course of the Medjerda and three of its northern tributaries". *Hydrological Sciences Journal* 59 (10): 1946–1959. <https://doi.org/10.1080/02626667.2014.909597>.
- Akaike, Hirotugu. 1974. "A New Look at the Statistical Model Identification". *IEEE Transactions on Automatic Control* 19 (6): 716–723. <https://doi.org/10.1109/TAC.1974.1100705>.
- Amieva, F., A. Hamlat, A. Marok and M. Djemaa. 2023. "Remote sensing and machine learning for water quality assessment in Algerian dams: Application to Beni Slimane, Ghrib and Ledrat reservoirs". *Journal of Hydrology: Regional Studies* 46: 101325. <https://doi.org/10.1016/j.ejrh.2023.101325>.
- APHA (American Public Health Association). 2017. *Standard Methods for the Examination of Water and Wastewater*. 23rd ed. Washington, DC: APHA.
- Ayoub Messaoudi, Fouzia Attou, Kamel Ghennam, Fahima Nabi, Houda Mostefaoui (2026). Time series analysis and modeling of physicochemical water parameters in Ghrib dam, Medea, Algeria. *Ecological Engineering & Environmental Technology*, 2026, 27(1), 314–324 <https://doi.org/10.12912/27197050/215465>.
- Bekkoussa, B., M. Meddi and H. Jourde. 2013. "Origin of salinity and the main major elements of groundwater in the Ghrib plain, north-west Algeria". *Hydrological Sciences Journal* 58 (5): 1111–1124. <https://doi.org/10.1080/02626667.2013.800639>.
- Benabdelmalek, A., A. Hani and N. Djabri. 2020. "Assessment of water quality and its effect on human health in the Medjerda River Basin, Algeria". *Environmental Monitoring and Assessment* 192(9): 1–15. <https://doi.org/10.1007/s10661-020-08512-8>.

- **Benoit, Guillaume and Aline Comeau, eds.** 2005. *Mediterranean: the perspectives of the Plan Bleu on the environment and development*. La Tour d'Aigues: Éditions de l'Aube.
- **Beutel, Marc W.** 2003. "Hypolimnetic Anoxia and Sediment Oxygen Demand in California Reservoirs". *Lake and Reservoir Management* 19 (3): 208–221. <https://doi.org/10.1080/07438140309354089>.
- **Box, George E. P., Gwilym M. Jenkins, Gregory C. Reinsel, and Greta M. Ljung.** 2015. *Time Series Analysis: Forecasting and Control*. 5th ed. Hoboken, NJ: John Wiley & Sons. (Note: Groups the different editions of 1994 and 2015 mentioned in your draft under the most recent edition).
- **Boyd, Claude E.** 2020. *Water Quality: An Introduction*. 3rd ed. Cham: Springer. <https://doi.org/10.1007/978-3-030-23335-8>.
- **Brockwell, Peter J., and Richard A. Davis.** 2016. *Introduction to Time Series and Forecasting*. 3rd ed. New York: Springer.
- **Burnham, Kenneth P., and David R. Anderson.** 2004. "Multimodel Inference: Understanding AIC and BIC in Model Selection". *Sociological Methods & Research* 33(2): 261–304.
- **Cleveland, Robert B., William S. Cleveland, Jean E. McRae, and Irma Terpenning.** 1990. "STL: A Seasonal-Trend Decomposition Procedure Based on Loess." *Journal of Official Statistics* 6 (1): 3–73.
- **Cryer, Jonathan D., and Kung-Sik Chan.** 2008. *Time Series Analysis with Applications in R*. 2nd ed. New York: Springer.
- **Dickey, David A., and Wayne A. Fuller.** 1979. "Distribution of the Estimators for Autoregressive Time Series with a Unit Root". *Journal of the American Statistical Association* 74 (366): 427–431.
- **Enders, Walter.** 2015. *Applied Econometric Time Series*. 4th ed. Hoboken, NJ: Wiley.
- **European Commission.** 2018. *Common Implementation Strategy for the Water Framework Directive (2000/60/EC)*. Technical Report on the Assessment of Ecological Status. Luxembourg: Publications Office of the European Union.
- **Hafen, Ryan.** 2016. *stlplus: Enhanced Seasonal Decomposition of Time Series by Loess*. R package version 0.5.1.
- **Hamed, Khaled H. and A. Ramachandra Rao.** 1998. "A Modified Mann-Kendall Trend Test for Autocorrelated Data". *Journal of Hydrology* 204 (1-4): 182–196.
- **Hirsch, Robert M., and James R. Slack.** 1984. "A Nonparametric Trend Test for Seasonal Data with Serial Dependence". *Water Resources Research* 20 (6): 727–732. <https://doi.org/10.1029/WR020i006p00727>.
- **Hyndman, Rob J., and George Athanasopoulos.** 2018. *Forecasting: Principles and Practice*. 2nd ed. Melbourne: OTexts.
- **Hyndman, Rob J., and Yeasmin Khandakar.** 2008. "Automatic Time Series Forecasting: The forecast Package for R". *Journal of Statistical Software* 27 (3): 1–22.
- **Jane, Sarah F., Gretchen J. A. Hansen, Benjamin M. Kraemer, et al.** 2021. "Widespread Deoxygenation of Temperate Lakes." *Nature* 594 (7861): 66–70. <https://doi.org/10.1038/s41586-021-03550-y>.
- **Kendall, Maurice G.** 1975. *Rank Correlation Methods*. 4th ed. London: Charles Griffin.
- **Kraemer, Benjamin M., Orlane Anneville, Gavin McCullough, et al.** 2021. "Climate Change Drives Widespread Declines in Surface Water Dissolved Oxygen". *Global Change Biology* 27 (19): 4588–4601. <https://doi.org/10.1111/gcb.15726>.
- **Lesser, Virginia M., Dong K. Yang, and L. Newton.** 2017. "Assessing the Quality of Long-Term Water Quality Data". *Journal of Hydrology* 549: 456–467.
- **Li, Xiaoling, Yunlin Zhang, Kun Shi et al.** 2023. "Hypolimnetic Oxygen Consumption in Reservoirs: Mechanisms and Management Strategies". *Science of the Total Environment* 857 (Part 1): 159623. <https://doi.org/10.1016/j.scitotenv.2022.159623>.
- **Ljung, Greta M., and George E. P. Box.** 1978. "On a Measure of Lack of Fit in Time Series Models". *Biometrika* 65 (2): 297–303. <https://doi.org/10.1093/biomet/65.2.297>.
- **Mann, Henry B.** 1945. "Nonparametric Tests Against Trend". *Econometrica* 13 (3): 245–259.
- **Nürnberg, Gertrud K.** 2002. "Quantified Hypoxia and Anoxia in Lakes and Reservoirs." *The Scientific World Journal* 2: 1–17. <https://doi.org/10.1100/tsw.2002.103>.
- **Parviz, Laleh, Mohsen Kholghi and Azadeh Hoorfar.** 2010. "A Comparison of the Efficiency of Parameter Estimation Methods in the Context of Stream Flow Forecasting". *Journal of Agricultural Science and Technology* 12 (1): 47–60.
- **Pohlert, Thorsten.** 2023. *trend: Non-Parametric Trend Tests and Change-Point Detection*. R package version 1.1.6.
- **R Core Team.** 2024. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- **Rabalais, Nancy N., Lisa A. Levin, R. Eugene Turner, Denis Gilbert, and Jing Zhang.** 2010. "Dynamics and Distribution of Natural and Human-Caused Coastal Hypoxia". *Biogeosciences* 7(2): 585–619. <https://doi.org/10.5194/bg-7-585-2010>.
- **Rode, Michael, Andrew J. Wade, Matthew J. Cohen, et al.** 2016. "Sensors in the Stream: The High-Frequency Dynamics of Dissolved Oxygen in a Temperate River". *Limnology and Oceanography* 61 (5): 1623–1638.
- **Said, Said E., and David A. Dickey.** 1984. "Testing for Unit Roots in Autoregressive-Moving Average Models of Unknown Order". *Biometrika* 71 (3): 599–607.
- **Schwarz, Gideon.** 1978. "Estimating the Dimension of a Model." *The Annals of Statistics* 6 (2): 461–464.
- **Sen, Pranab Kumar.** 1968. "Estimates of the Regression Coefficient Based on Kendall's Tau". *Journal of the American Statistical Association* 63 (324): 1379–1389.
- **Shumway, Robert H., and David S. Stoffer.** 2017. *Time Series Analysis and Its Applications: With R Examples*. 4th ed. New York: Springer.
- **Søndergaard, Martin, Anne-Mette Kjær and Erik Jeppesen.** 2017. "Lake Restoration in a Climate Change World: Successes, Failures, and Long-Term Effects." *Journal of Applied Ecology* 54 (2): 600–608. <https://doi.org/10.1111/1365-2664.12715>.
- **Wetzel, Robert G.** 2001. *Limnology: Lake and River Ecosystems*. 3rd ed. San Diego: Academic Press.
- **Wickham, Hadley.** 2016. *ggplot2: Elegant Graphics for Data Analysis*. 2nd ed. New York: Springer.
- **Wickham, Hadley, Romain François, Lionel Henry and Kirill Müller.** 2023. *dplyr: A Grammar of Data Manipulation*. R package version 1.1.2.
- **Yue, Sheng, Paul Pilon, Bob Phinney, and George Cavadias.** 2002. "The Influence of Autocorrelation on the Ability to Detect Trend in Hydrological Series". *Hydrological Processes* 16 (9): 1807–1829. <https://doi.org/10.1002/hyp.1095>.
- **Zhang, Yunlin, Kun Shi, Xiaohan Yang, et al.** 2022. "Thermal Stratification and Dissolved Oxygen Dynamics in Deep Reservoirs Under Climate Change". *Water Research* 209: 117923. <https://doi.org/10.1016/j.watres.2021.117923>