

Hyperspectral Remote Sensing of Floristic Composition and Phytogeographical Distribution in Gebel El-Sibaie, El-Quseir, Egypt

Nessrien m. Hamad-allah ^{*1,2}, Amal A. Morsy², El-Sayed A. Khalifa³, Ghada A. Khdery⁴, Mahmoud E. Ali⁵, Maged M. Abutaha⁶

^{1,2} Faculty of Science, Ain Shams University, Cairo, Egypt.

^{3, 5,6} Department of Plant Ecology and Range Management, Desert Research Center, Cairo, Egypt

⁴Department of Agriculture application, National Authority for Remote Sensing and Space Science, Cairo, Egypt

Corresponding author: Nessrien M. Hamad-Allah (nessrien2023@gmail.com)

NESSRIEN M. HAMAD-ALLAH (nessrien2023@gmail.com)

Amal A. Morsy (amalmorsy@sci.asu.edu.eg)

EL-SAYED A. KHALIFA (nessoali2025@gmail.com)

GHADA A. KHDERY (ghada.ali@narss.sci.eg)

MAHMOUD E. ALI (mdsw78@hotmail.com)

MAGED M. ABUTAHA (maged_abutaha@yahoo.com)

Abstract

Our study aims to explore the flora of the Gebel El-Sibaie and utilizes hyperspectral remote sensing to enhance the assessment of plant diversity. We analyze vegetation elements, including floristic diversity, life form, lifespan, and phytogeographical affinities, and utilize hyperspectral remote sensing with an ASD spectroradiometer. A total of 60 species belong to 49 genera and 29 different families, with Fabaceae, Zygophyllaceae, and Asteraceae being the most characteristic families. The flora exhibits high monotypism, with 40 genera and 19 families. Chamaephytes and therophytes are the most prevalent life forms. The common lifespan was for perennials. *Acacia cyclocarpa*, *Leucaena leucocephala*, and *Prosopis juliflora* were recorded as alien species for the first time in Al-Quseir. Phytogeographical analysis reveals that mono-regional elements are the most dominant chorotypes, with Saharo-Arabian taxa comprising approximately a quarter of the floristic structure. The study created a spectral library of 37 plants and soil samples from 22 sites. It found consistent spectral signatures across plants, especially in the visible and infrared regions. *Phragmites australis*, *Leucaena leucocephala*, and *Paronychia argentea* had higher reflectance. Furthermore, plants manage water stress differently, with higher visible reflectance associated with increased chlorophyll content and lower infrared reflectance resulting from water absorption. Soil properties like coarse sand enhance reflectance in the visible and NIR regions, while silt and clay boost absorption in the SWIR. Chemical factors, such as EC and mineral content, impacted spectral signatures.

Keywords: Arid mountain, Chorology, Diversity, invasive, life form, Red Sea, Spectral Signature

1. Introduction

The Eastern Desert of Egypt, extending eastward from the Nile Valley to the Red Sea and the Gulf of Suez, comprises 21% of Egypt's total land area. Moreover, comprises the backbone of high, rugged mountains parallel to the Red Sea coast. Its major phytogeographical regions are the Red Sea coastal region and the inland Sahara (Zahran & Willis, 2009). The Egyptian

desert, with its arid and/or extremely arid climate, exposes vegetation to harsh and drastic environmental circumstances (Batanony, 1979; Zahran & Willis, 2009).

Desert vegetation was classified into three categories: a permanent framework of perennials, ephemerals, and accidental species, which are characterized by low and erratic rainfall, the likelihood of rainy seasons, and soil thickness, which affects the appearance and lifespan of ephemerals (Kassas, 1952, 1966, 1971). Ephemerals, annuals, and biennial plants become drier with the arrival of a lengthy summer rainless season (Boulos, 2005). Perennials, a common framework for desert plants and a good predictor of habitat conditions (Zahran & Willis, 2009), are classified into five groups: xerophytes, halophytes, psammophytes, helophytes, and mangroves (Zahran & El-Amier, 2014). Perennial xerophytes, drought-resistant shrubs, and under shrubs are the most significant and distinctive plant life in the Egyptian desert (Zahran & Willis, 2009).

Desert vegetation exhibits significant variation in species distribution with minimal irregularities in physiographic factors, making it an interesting and informative aspect of plant ecology (El-Khouly & Shawky, 2017). Vegetation structure, floristic composition, and life form spectrum are crucial for ecological studies. They are demonstrated through vertical species stratification, spatial distribution of individuals, and species abundance (Barnes et al., 1998; Abutaha et al., 2021).

Abiotic, biotic, historical, and human influence plant distribution and abundance (Wisiz et al., 2013). Irresponsible anthropogenic interventions, harsh habitat conditions, climate change, and alien species negatively impact desert wildlife population dynamics, threaten biodiversity, disrupt natural habitats, and rapidly deplete natural vegetation in arid regions (Hegazy et al., 2004a, 2007; Alhammadi, 2010; Halmy et al., 2019; Mahklouf, 2019; Bedair et al., 2021; Hussein et al., 2021; Abbas et al., 2023; Haq et al., 2024).

Egypt is divided into three floral provinces: (1) Saharo-Sindian Province “includes most of Egypt’s deserts”; (2) The Iranian Turanian “includes the mountainous region of South Sinai and the Galala Mountains of the Eastern Desert”; (3) The Sudano-Zambezian “includes the Gebel Elba region in southeastern Egypt” (Eig, 1931-1932). Most of Egypt's deserts are within the Sahara Regional Transition Zone, except for the Western Mediterranean coastal zone and the Gebel Elba region, which are part of the Sahel region (White, 1993).

The hyperspectral technique offers a range of absorption based on spectral narrow bands to distinguish between different plants based on their spectral signature (Nagendra, 2002). The reflectivity of plant features is associated with biological substances that are present in the leaves, and the physical attributes of the leaves, including the air-water interfaces and air-space fraction. Plant physiology, morphology, or anatomy provides the basis for a species' spectral characteristics (Jarocinska et al., 2016; Kycko et al., 2014). It is possible to compute, record, and assign even slight variations in reflectance to specific plants, enabling the classification of species based on their spectral characteristics. In this manner, species-specific spectral characteristics were identified using contrasting spectral reflectance patterns (Kupková et al., 2017; Marcinkowska et al., 2017). Khderi et al. (2014); Aboelghar and Khderi (2017); Khderi et al. (2019); Yones (2019); Gamal et al. (2020), and others are just a few examples of the various applications of multispectral and hyperspectral data in Egypt, including vegetation cover discrimination and vulnerability assessments.

The present study was designed to investigate the current status of the floristic composition in the Gebel El-Sibaia area, encompassing mountains and wadis distributed throughout the study area. It compares the present status of flora with earlier records and relevant studies, recognizing life forms and chorological distribution of the vegetation. The study utilized

hyperspectral remote sensing, employing an ASD spectroradiometer, to enhance the assessment of plant diversity and distribution patterns within this unique environment.

2. Materials and methods

2.1. Study area

The Study area is a part of the hyper-arid Eastern Desert of Egypt. It is situated nearly in the middle part of the Eastern Desert, South-West of Al-Quseir city by about 30 Km, and located between latitudes 25° 30' and 25° 45' N and longitudes 34° 0' and 34° 30' E. It consists of Gebel El-Sibaie, Gebel Umm lasifa, and Gebel Umm Naqqat, and their main drainage systems, from north to south, namely: Wadi Esel, Wadi Sharm El-Bahari, and Wadi Umm Gheig, leading to the coastal salt marshes along the Red Sea coast. (Fig. 1).

2.2. Floristic composition

The fieldwork was conducted during 2021, 2022, and 2023 through several trips. A total of 24 vegetation stands were randomly chosen, based on the diversity and variation of habitat types in the study area. For every stand, the species composition was listed. Plant species nomenclature and identification followed Täckholm (1974) and were updated according to Boulos (1999, 2000, 2002, 2005, 2009). The annual and perennial (life span) species were listed. The plant life forms are identified according to Raunkiaer (1934). The phytogeographical affinities of the surveyed species were defined as in previous studies on the Eastern Desert of Egypt (Salama et al., 2014, 2023; Abd El-Ghani et al., 2013, 2017; Amro et al., 2021).

2.3. Soil sampling and analysis

Mixed soil samples were collected from the surface layer (0-20 cm) for each stand, air-dried, sieved through a 2 mm sieve to remove gravel and debris, and then packed to be ready for analysis. Physical and chemical analyses of soil samples were carried out according to (Richards, 1954; Jackson, 1962; Krumbein & Sloss, 1963; Allen et al., 1974; Ryan et al., 1996; ASTM, 2002).

2.4. Spectroscopic Analyses

The reflectance of 37 plant leaves was measured in the field using the ASD Field Spec. Spectrum reflectance curves were used to visually verify the vegetation spectrum reflectance qualities after the data was saved to an ASCII file using the statistical analysis application ASD View Pro. A statistical analysis identified the spectral bands that best distinguish the plants under investigation. The selection of bands was carried out using linear discriminant analysis (LDA) (Clark et al., 2005), and the effectiveness of the chosen wavelengths and bands in differentiating plant species was examined using ANOVA (Anderson, 2001).

3. Results

3.1. Floristic Diversity

A total of 60 plant species belonged to 49 genera, and 29 different families were recorded in the total study area; they were made up of 42 perennials (70.00%) and 18 annuals (30%) (Appendices 1, Fig. 2). Among them three species (about 5.00 %) of the total surveyed flora have been recorded for the first time and represent new additions to the flora of Al-Quseir region; which are: *Acacia cyclocarapa*, *Leucaena leucocephala*, and *Prosopis juliflora*. These new records pertain to invasive taxa within the family Fabaceae (Appendix 1). Moreover, one species is considered endangered in Egypt (*Moringa peregrina*).

The species-rich families were Fabaceae (11 species, 18.33%), Zygophyllaceae (7 species, 11.67%), and Asteraceae (5 species, 8.33%) in the total study area, followed by Caryophyllaceae, represented by four species (6.67%). Meanwhile, Cleomaceae and Poaceae were represented by three species each (5.00%). Asclepiadaceae, Boraginaceae, Capparaceae, and Resedaceae exhibited the same number of recorded species (2 species=3.33% each). On the other hand, 18 families were poorly represented, having only one species each (1.67%) (Fig. 3). The most characteristic families in the study locations were Fabaceae in both Wadi bed of Um Gheig (18.75%) and Esel (26.92%), and Zygophyllaceae at G. El-Sibaie, G. Um-Nakkat, and Wadi bed of Sharm-El-Bahary (18.52%, 13.79% and 25% respectively). Gebel Umm Lasifa is the lowest location, comprising four families, each with one species (25%), namely Resedaceae, Moringaceae, Fabaceae, and Capparaceae (Appendix 1).

The most prominent families in terms of the number of genera in the total study area were Fabaceae (7 genera), Asteraceae, Caryophyllaceae & Zygophyllaceae (4 genera each) and Poaceae (3 genera). Meanwhile, the genera with the most significant number of species were *Acacia* L. with four species (6.67%), *Cleome* and *Zygophyllum* with three species each (5.00%), *Pulicaria*, *Capparis*, *Astragalus*, and *Fagonia* with two species each (3.33%). On the other hand, the other genera comprise the lowest number, one species each (1.67%).

The study clearly showed that the slope of mountains support vegetation growth of about 36 plant species and is dominated mainly by *Acacia tortilis*, *Capparis cartilaginea*, *Cleome droserifolia*, *Leptadenia pyrotechnica*, *Lycium shawii*, *Moringa peregrina*, *Ochradenus baccatus*, *Reseda pruinosa*, *Tephroisa apollinaea*, and *Zygophyllum simplex*. Open desert is characterized by 41 species, and is dominated mainly by *Zilla spinosa*, *Zygophyllum coccineum*, *Acacia tortilis*, *Balanites aegyptiaca*, *Capparis cartilaginea*, *Cleome droserifolia*, *Reseda pruinosa* (Appendix1). Mangrove vegetation which is characterized by only one species *Avicennia marina*, and salt marsh vegetation including about 14 species *Nitraria retusa*, *Limonium axillare*, *Phragmites australis*, *Salsola imbricata*, *Tamarix nilotica*, *Taverniera aegyptiaca*, and *Zygophyllum album*. Overall, Gebel Umm Lasifa was the least diverse among the studied area.

3.2. Life form

Fig. 4 depicts five different life forms of the 60 plant species recording in the study area. Chamaephytes (Ch) & Therophytes (Th) were the dominant life forms each represented by 30.00% of the total number of flora, most common species were *Zilla spinosa* (L.), *Capparis sinaica* Veil., *Cleome droserifolia* (Forssk.), *Fagonia mollis* Delile *Zygophyllum coccineum* L & *Cometes abyssinica* R.Br., *Reseda pruinosa* Delile, *Zygophyllum simplex* L. *Tribulus terrestris* L and *Aizoon canariense* L. Followed by Phanerophytes (Ph) which comprises around 28.33% of the recorded species, of which *Avicennia marina* (Forssk.), *Leptadenia pyrotechnica* (Forssk.), *Acacia tortilis* subsp. *raddiana* (Savi), *Acacia tortilis* (Forssk.), *Lycium shawii* Roem., *Moringa peregrina*, *Ochradenus baccatus*. Hemicryptophytes account for approximately 10.00% of the vegetation and included 6 species *Trichodesma ehrenbergii* Schweinf, *Phragmites australis* (Cav.), *Citrullus colocynthis* (L.), *Kickxia acerbiana*(Boiss.), *Forsskaolea tenacissima* L. and *Polycarpaea repens* (Forssk.) On the other hand, the lowest life form recorded is Geophytes (1.67 %) represented only by one species (*Panicum turgidum* Forssk.).

Fig.5 shows that Phanerophytes and chamaephytes were the most represented life forms in all study sites, with a high presence of (30.23 and 24.44%) at Wadi Esel and G. Umm-Naqqat, respectively, and a low presence (4.65 and 2.22% respectively) at Wadi Sharm El-Bahary and G. Umm Lasifa (no other life forms have been recorded at this location). Therophytes prevail (44.83%) at Wadi Umm-Gheig, while they are absent from Wadi Sharm El-Bahary and G.

Umm-Lasifa. The highest value of Hemicryptophytes (42.86%) was recorded at El-Sebaie Mountain. On the other hand, Geophytes were reported only at Wadi Umm-Gheig and Umm Naqqat Mountain with equal percentage (50% each).

3.3. Chorology

According to phytogeographical affinities, the total recorded species were classified into three major phytogeographical groups: monoregional, bioregional and pluriregional, in addition to, pantropical and Afro-Montane species (Table 1 & Fig. 6). Monoregional elements were the highest (29 species = 48.33%) among the surveyed species in the study area with different affinities, and fall under five main chorotypes: Saharo-Arabian had a wide geographical range (15 species accounting 25.00%); Saharo-Sindian (6 species forming 10.00%); Sudano-Zambezian (3 species forming 5.00%); the last two chorotypes were Sudanian & Medeterranean represented only by one species (*Acacia ehrenbergiana* Hayne & *Paronychia argentea* Lam.) each forming 1.67 %. Two American taxa, *Leucaena leucocephala* and *Prosopis juliflora*, represent 3.33%. One Australian taxon (*Acacia cyclocarpa*), representing 1.67%.

Twenty-two species of all recorded species (36.76%) were biregional taxa, and fall under four chorotypes: the Saharo-Arabian/Sudano-Zambezian chorotypes together had the highest share of 13 species (21.67%), Saharo-Arabian/Irano-Turanian regions, represented by seven species (11.67%). At the same time, Saharo-Arabian/Sudanian & Mediterranean/Saharo-Arabian regions have only one species (1.67% each), *Capparis decidua*, and *Phoenix dactylifera* L., respectively. The pluriregional elements were represented by a total of 6 species (10.00%) of different affinities, which fall under five main chorotypes: Mediterranean/Saharo-Arabian/Irano-Turanian, with the highest share (2 species, forming 3.33%). At the same time, the other four chorotypes are Mediterranean/Saharo-Arabian/Sudano-Zambezian, Mediterranean/Saharo-Sindian/Irano-Turanian, Mediterranean/Irano-Turanian/Euro-Siberian, and Saharo-Arabian/Sudano-Zambezian/Irano-Turanian. Only one species each (forming 1.67% of recorded species). The three remaining species were distributed among two species (3.33%) of the Pantropical chorotype (*Phragmites australis* (Cav.) and *Tribulus terrestris* L.) and one species of the Afro-Montane (*Cometes abyssinica* R. Br.), forming 1.67%.

3.4. Soil properties

Physical and chemical analysis of the soil of the study area were represented in Appendix(2,3). The results of granulometric analysis showed that the soil texture was primarily defined by a high percentage of very coarse and coarse sand at the slopes of the mountains, i.e., Gebel Umm Nakkat, Umm Lasifa, and El-Sibaie. Open deserts were characterized by delicate and very fine sand with high percentage at wadi Esel (24.43, 45.85 respectively). At the same time, mangrove habitat was characterized by high percentage of Silt & Clay (41.44) at Wadi Sharm El-Bahary, followed by salt marshes habitat (37.69 %) at wadi Esel. For chemical analysis, the soils of the study area were characterized as slightly acidic to alkaline, with the mean pH value ranging from 6.26 to 9.00 (Appendix 3). The EC ranges between (0.20–25.70 mS/m). The mangrove habitat recorded the highest EC (25.68), followed by halophytic habitat, and wadi bed of wadi Umm Gheig (9.50, 6.69, respectively), while the slope of Gebel El-Sibaie records the lowest EC (0.22). TDS, and most mineral contents; i.e., Ca⁺⁺, Mg⁺⁺, Na⁺, HCO₃⁻, SO₄²⁻, and Cl⁻ were high at Wadi Sharm El-Bahary. In contrast, P record the highest value at wadi bed of Gebel El-Sebaie.

3.5. Hyperspectral analysis and application

3.5.1 Hyperspectral signature for plants

Fig. 7 illustrates the typical spectral signatures of 37 plant species that are found naturally in the study area. The results revealed a high degree of consistency in the spectral signatures across different plants, particularly in the visible and infrared regions of the spectrum. Among all spectral regions, *Phragmites australis*, *Leucaena leucocephala*, and *Paronychia argentea* showed higher reflectance compared to other species. *Avicennia marina* exhibited the lowest reflectance in the SWIR 1 and SWIR 2 zones (Fig.8). Additionally, *Leucaena leucocephala* and *Salsola imbricata* recorded higher reflectance values in the visible region than other species (Fig.8).

3.5.2 Analysis of Soil Reflectance Data

Fig. 9 illustrates the typical spectral signatures of soil samples from 22 different sites, showing variability across the visible (350-700 nm), near-infrared (NIR, 700-1400 nm), and shortwave-infrared (SWIR, 1400-2500 nm) regions (Fig. 10). In the visible Region (350-700 nm), reflectance generally increases with wavelength. Soil sites 4, 5, and 7 show higher reflectance, indicating lower organic matter and higher mineral content, while in the Near-Infrared Region (700-1400 nm), High reflectance values suggest a high content of quartz and other silicate minerals. Soil sites 4, 5, and 11 exhibit higher reflectance, indicating robust mineral composition. In SWIR 1 & 2 (1400-2500 nm), multiple absorption features correspond to water and clay minerals (Fig. 10). Soil sites 7 and 16 show pronounced absorption, indicating higher clay and moisture content.

4. DISCUSSION

Floristic studies are not only important for exploring the floral diversity of an area, but are also of social and economic importance (Al-Mutairi et al. 2016). Floristic studies, despite being traditional, are crucial in modern botanical research as they help slow down human pressure on biodiversity, manage actions, and maintain conservation strategies (Wagensommer et al. 2023). The wild vegetation in the study area is composed of 42 perennials (70.00%) and 18 annuals species (30.00%) of the total recorded species. The current study's prevalence of perennials agrees with the findings of (Zahran & Willis. 2009; Abdelaal et al. 2016 and Bedair et al. 2021 on Wadi Hagul) and most recently (Salama et al. 2023) in the Eastern Desert of Egypt. The vegetation structure of some Wadis at the Red Sea coast is simple, requiring perennial species to endure harsh environmental conditions (Shawky and El-Khouly 2017). The desert harsh conditions pose a significant risk to the survival of plant species, particularly annuals (Hegazy et al., 2008).

The study identifies three key families as leading taxa that constitute the major bulk flora of the study area were Fabaceae (18.33%), Zygophyllaceae (11.67 %) and Asteraceae & (8.33 %), which contributed collectively about 38.33 % of the total recorded species. Comparable results have been reported in other studies (e.g., Abd El-Ghani et al., 2013; Kurd et al., 2017; El-Khouly and Shawky, 2017 and Salama et al., 2018). Amro et al. (2021) demonstrated that the most prevalent flora in the North African and Mediterranean are Asteraceae, Zygophyllaceae, Fabaceae, Chenopodiaceae and Brassicaceae. Salama et al. (2023) reported that the four families Asteraceae, Poaceae, Fabaceae, and Zygophyllaceae make up the majority of the flora in the eastern desert of Egypt.

Asteraceae, Poaceae and Fabaceae represent the majority of Egyptian flora and adjacent countries such as Libya and Saudi Arabia (Täckholm 1974; Boulos 2009; El-Barasi et al.,

2011 and Ahmed et al. 2020). The highest among other perennial species were *Zilla spinosa* and *Zygophyllum coccineum* in the studied stands and spread their dominance all over the Eastern Desert of Egypt (Shaltout et al. 2009; Abd El-Ghani et al. 2013a; Amro et al. 2021 and Salama et al. 2023).

According to El-Behairy et al. (2020) 41 alien species were not recorded in the last updated checklist of the Egyptian flora (Boulos, 2009). Among these invasive species is *Prosopis juliflora*, which spread clearly and has a negative impact on the biodiversity in the southeastern corner of Egypt (Mohamed et al., 2015 and Slaltout et al., 2016). It reduced plant diversity (Allelopathic effect), because it has massive impacts upon water resources (Weber, 2003), represent a tangible threat to the sustainability of existing natural resources in range and cultivated lands (Assaeed et al., 2020; Badawy et al., 2021 and Bedair, et al. 2021).

Kassas (1952) pointed out that Life form distribution of the desert plants is closely correlated with rainfall, topography and landform. The life form evolved as a result of climatic and environmental adaptation (Hegazy et al., 2012 and Salama et al., 2019). The present study demonstrated that different plant species were categorized to five life forms, Chamaephytes & therophytes (30.00 % each) and Phanerophytes (28.81%) constituted the largest number of species (53) with total value (88.33 %). While Hemicryptophytes (10.00%), Geophytes the lowest one (1.67%), are represented by only one species. Our results agreed with other reports (Abd El-Ghani et al., 2000 & 2013; Kurd et al., 2017; Shawky and El-Khouly 2017 and Salama et al., 2023) in the Eastern desert. According to (Hussien et al. 2019), chamaephytes were the most abundant life form, followed by therophytes, hemicryptophytes and phanerophytes in hyper-arid desert environments. From phanerophytes species, *Moringa peregrina* was the most endangered tree where there is a sharp decline in its numbers and sizes of its populations in Egypt (Abd El-Wahab, 1995; Hegazy et al., 2007; Dadamouny, 2009 and Zaghloul et al., 2012).

Another aspect of floristic research is chorological affinity, which pertains to regional distributions and plant responses to environmental changes (Salama et al., 2023). Ecological adaptations of species affect how widely distributed a plant is (Zahran, 1982; Zahran and Willis 2009 and Hammad et al., 2024). For example, mangrove community occur in a zone following the shore-line, while other vegetation zones may include a mosaic of one or more than one community depending on local topography or soil conditions (Zahran and Willis, 2009).

The chorological analysis of the surveyed flora revealed that 29 species (48.33% of the total flora) were monoregional, of which 15 species (25.00%) were pure Saharo-Arabian chorotype. Saharo-Arabian species are abundant in the eastern desert in Egypt, this is may be attributed to their high tolerability to harsh desert environmental conditions (Amro, et al., 2021 AND Salama, 2023). 3 species (5.00%) included *Aizoon canariense*, and *Moringa peregrina* represented the typical (mono-regional) Sudano Zambeziian chorotype, the same results reported by (Amro, et al., 2021). The bioregional chorotype ranked second (36.67%), of which Saharo-Arabian, Sudano-Zambeziian element have the highest share of the recorded species (21.67%). On the other hand, pluri-regional with low share (10.00%) and extending their distribution all over the Saharo-Arabian, Sudano- Zambeziian, Irano-Turanian and Mediterranean regions. Our results coincide with others obtained by (Abutaha et al., 2019, 2020; Amro et al., 2021 and Salama et al., 2023) in Eastern desert in Egypt and (Ahmed et al., 2020) in Saudi Arabian.

The spectral signatures of plant species provide insights into their ecological adaptations and health. For instance, species like *Phragmites australis* and *Leucaena leucocephala* show higher reflectance in various spectral regions, reflecting their robust physiological conditions and ability to thrive in diverse environments. Dominant life forms such as chamaephytes (low-growing perennials) and therophytes (annuals) have distinct spectral characteristics linked to their photosynthetic activity and strategies for surviving harsh conditions. The study area predominantly features mono regional species, particularly Saharo-Arabian elements, adapted to arid environments. These species exhibit spectral signatures indicative of efficient water use and drought resistance. The spectral data reveal how different species manage water stress, with certain plants showing higher reflectance in the visible region due to chlorophyll content and others demonstrating lower reflectance in the infrared region due to water absorption. This connection between spectral signatures and ecological traits helps in understanding the health, stress levels, and adaptive strategies of the plant species in their natural habitats (Khderiy 2019, 2020, 2021, 2023, Aboelghar and Gamal et al., 2020 and El-Shirbeny, 2021).

The results demonstrate that soil texture, chemical composition, and moisture content are key factors affecting the spectral behavior of soil. Coarse-grained soils with low organic matter tend to have higher reflectance across the visible and NIR regions due to their mineral composition. In contrast, fine-grained soils with higher silt and clay content exhibit absorption features in the SWIR range, which are linked to water and clay minerals. Coarse sands found on the slopes of Gebel Umm Nakkat, Umm Lasifa, and El-Sibaie reflect more light in the visible spectrum due to lower organic content and larger particle size. Fine sands, particularly in Wadi Esel, scatter light and show moderate reflectance in the NIR region, where quartz and silicate minerals dominate. Mangrove habitats, with high silt and clay content (e.g., Wadi Sharm El-Bahary), show strong absorption in the SWIR range due to water and clay minerals.

Soils with a pH of 6.26–9.00, especially clay-rich soils, display absorption in the SWIR region, indicating moisture interaction. High electrical conductivity (EC) in mangrove and halophytic soils results in lower reflectance, while low-EC soils, such as Gebel El-Sibaie, exhibit higher reflectance in the NIR. Sites rich in quartz and silicates, like Gebel El-Sibaie, reflect strongly in the NIR range, while high mineral content (e.g., calcium and magnesium) in Wadi Sharm El-Bahary influences the spectral signature in the SWIR range through water and mineral absorption.

Conclusion

The present work is the first floristic study of Gebel El-Sebaie area, we found that vegetation within study area is restricted to a definite habitat depending on its ability to adapt to drought stress. Soil physical and chemical properties strongly influence their spectral signatures, supporting the use of remote sensing to differentiate soil types and environmental conditions. Our study is attempting to cover the lack of and /or insufficient knowledge of plant diversity of less investigated regions, such as the Red Sea Mountain.

Author contributions

M.E. A., M. M. A., and N.M.H. carried out fieldwork, identification of species, and Conceptualization. Gh. A. Kh., M. M. A., and N. M. H. wrote original draft preparation, methodology, software, data curation, and formal analysis., M. M. A. and A. A. M. reviewed and edited the draft, while all authors contributed to the final version.

Acknowledgements

I would like to extend my sincere appreciation to my colleagues at the Desert Research Center, Egypt, Dr. Ahmed Mandouh and Dr. Mohamed Kashlana, for their support during field trips and the collection of plants and soil.

Funding Information

This research received no external funding.

Data Availability Statement

Data underlying the findings of this study will be made available from the corresponding author upon reasonable request.

REFERENCES

- [1] Abbas AM, Soliman WS, Alomran MM, Alotaibi NM, Novak SJ (2023) Four Invasive Plant Species in Southwest Saudi Arabia Have Variable Effects on Soil Dynamics. *Plants*, 12, 1231. <https://doi.org/10.3390/plants12061231>
- [2] Abd El-Ghani MM (2000) Floristics and environmental relations in two extreme desert zones of western Egypt. *Global Ecol Biogeogr*, 9: 499-516. Doi:10.1046/j.13652699.2000.00209.x
- [3] Abd El-Ghani MM, Salama F, Salem B, El-Hadidy AM Abdel-Aleem M (2013a) Biogeographical relations of a hyperarid desert flora in eastern Egypt. *Afr. J. Ecol.*, 52:173-191. <http://dx.doi.org/10.1111/aje.12102>
- [4] Abd El-Ghani MM, Salama F, Salem B, El-Hadidy AM Abdel-Aleem M (2017). Phytogeography of the Eastern Desert flora of Egypt. *Wulfenia*, 24: 97-120.
- [5] Abd El-Wahab RH (1995) Reproduction Ecology of Wild Trees and Shrubs in Southern Sinai, Egypt. M.Sc. Thesis, Botany Department, Faculty of Science, Suez Canal University, Ismailia, Egypt. 110 p.
- [6] Aboelghar M, Khder G (2017) Spectroscopic and morphological characteristics of genus *Jatropha* (Euphorbiaceae) and genus *Jojoba* (Simmondsiaceae). *Asian J. Agric. & Biol.*, 5: 47-50.
- [7] Abutaha MM, El-Khouly AA, Jurgens N, Morsy AA, Oldeland J (2019) Elevation-richness pattern of vascular plants in wadis of the arid mountain Gebel Elba, Egypt. *Afr J. Ecol.*, 57(2): 238-246. <https://doi.org/10.1111/aje.12593>
- [8] Abutaha MM, El-Khouly AA, Jurgens N, Oldeland J (2020) Plant communities and their environmental drivers on an arid mountain, Gebel Elba, Egypt. *Vegetation Classification and Survey*, 1: 21-36. <https://doi.org/10.3897/VCS/2020/38644>
- [9] Abutaha MM, El-Khouly AA, Jurgens N, Oldeland J (2021) Predictive mapping of plant diversity in an arid mountain environment (Gebel Elba, Egypt). *Appl Veget Sci.*, 24(2), e12582. <https://doi.org/10.1111/avsc.12582>
- [10] Ahmed MA, Mohammed AK, Mohammad YA, Serag EE, Mohamed OB (2020) Floristic Diversity and Phytogeography of JABAL Fayfa: A Subtropical Dry Zone, South-West Saudi Arabia. *Diversity*, 12:1-20. <https://doi.org/10.3390/d12090345>
- [11] Al-Gifri AN, Kasem WT, Shehata RS, Eldemerdash MM (2019) The African Paleotropical Influence on the Biogeography of the Flora of Jazan, KSA. *Asian J. Soil Sci. Plant Nutr*, 4: 1-10. DOI: 10.9734/AJSSPN/2019/46160
- [12] Alhammadi ASA (2010) Preliminary survey of exotic invasive plants in some western and high plateau mountains in Yemen. *Ass. Univ. Bull. Environ. Res.*, 13 (1): 1-14. Doi: 10.21608/auer.2010.149259
- [13] Allen SE, Grimshaw HM, Parkinson JA, Quarmby C (1974) Chemical analysis of Ecological materials. Blackwell Scientific publication. Oxford; 565 p.
- [14] Amro A, Salama FM, Abd El-Ghani MM, El-Zohary AM, El-Shazoly RM (2021) Variations in community structure and plant species diversity with soil properties in a hyper-arid coastal desert of Egypt. *J. of Anim. Plant Scie.*, 31:1686-1698. DOI:10.36899/JAPS.2021.6.0372
- [15] Anderson JE, Kono TJ, Stupar RM, Kantar MB and Morrell PL (2016) Environmental association analyses identify candidates for abiotic stress tolerance in Glycine soja, the wild progenitor of cultivated soybeans. *G3 (Bethesda)* 6 (4): 835-843. DOI: 10.1534/g3.116.026914
- [16] ASTM (American Society for Testing and Materials) (2002) Annual book of ASTM standards. Baltimore, MD, USA, Vol. 11.01, 939 pp.
- [17] Barnes BV, Zak DR, Denton SR, Spurr SH (1998) *Forest Ecology*, fourth ed. John Wiley and Sons, New York.

- [18] Batanouny KH (1979) Vegetation along Jeddah– Mecca road: pattern and processes as affected by human impact. *J. Arid Envir.*, 2: 21–30. DOI:10.1016/S0140-1963(18)317014
- [19] Bedair R, Ibrahim AA, Alyamani AA, Aloufi S, Ramadan S (2021) Impacts of Anthropogenic Disturbance on Vegetation Dynamics: A Case Study of Wadi Hagul, Eastern Desert, Egypt. *Plants*, 10, 1906. <https://doi.org/10.3390/plants10091906>
- [20] Boulos L (1999) 'Flora of Egypt.' Vol. I (Azollaceae –Oxalidaceae). Al hadara Publishing, Cairo, Egypt, 419 pp.
- [21] Boulos L (2000) Flora of Egypt. Vol. II (Geraniaceae -Boraginaceae). Al Hadara publishing, Cairo, Egypt, 352 pp.
- [22] Boulos (2002) Flora of Egypt. Vol. III (Verbenaceae -Compositae). Al Hadara publishing, Cairo, Egypt, 373 pp.
- [23] Boulos L (2005) Flora of Egypt. Vol. IV (Alismataceae -Orchidaceae). Al Hadara publishing, Cairo, Egypt, 617 pp.
- [24] Boulos L (2009) Flora of Egypt-Checklist 2nd ed. Revised annotated edition. Al Hadara publisher, Cairo, Egypt. 2009;292 p.
- [25] Clark ML, Roberts DA, Clark DB (2005) Hyperspectral discrimination of tropical rain forest tree species at leaf to crown scales. *Remote Sens. Environ.*, 96 (3–4): 375–398. <https://doi.org/10.1016/j.rse.2005.03.009>
- [26] Dadamouny M (2009) Population Ecology of *M. peregrina* growing in Southern Sinai, Egypt. M.Sc. thesis, Botany Department, Faculty of Science, Suez Canal University, Ismailia, Egypt. DOI:10.13140/RG.2.1.5091.9760
- [27] Eig A (1931–1932) Les elements et les groupes phytogeographiques auxiliaires dans la flore palestinienne. *Beih. Repert. Spec. Nov. Regni. Veg.*, 63 (1, texte): 1–201 (1931); (2, Tabl. analyt.), 1–120 (1932).
- [28] El-Barasi Y M, Saeed M W B, El-Amrouni Abdelsalam O, Mohamed NF (2011) Check list of flora and vegetation on south elmarj zone: South EL-JABAL EL-Akhadar – Libya. *International J. of Engineering*, 3: 141-146.
- [29] El-Shirbeny MA, Ali AM, Khdery GA, Saleh NH, Afify NM, Badr MA, Baioumy EM (2021) Monitoring agricultural water in the desert environment of New Valley Governorate for sustainable agricultural development: a case study of Kharga. *Euro-Mediterranean J. Envi. Integration.*, 6 (2): 56,1-15. DOI:10.1007/s41207-021-00256-5
- [30] El-Zeiny A, Khdery G, Gad Abd-Alla (2022) Hyperspectral based approach to investigate topsoil characteristics of different taxonomic units of El-Fayoum depression. *Egypt. J. Remote Sensing Space Sci.*, 25: 405-415. <https://doi.org/10.1016/j.ejrs.2022.02.007>
- [31] Gamal E, Khdery G, Morsy A, Ali A, Ahmed Hashim A, Saleh S (2020) Using GIS based modeling to aid conservation of two endangered plant species (*Ebenus Armitagei* and *Periploca Angustifolia*) at Wadi Al-Afreet, Egypt. *Remote Sens. Applications: Society Envi.*, 19 100336. <https://doi.org/10.1016/j.rsase.2020.100336> 2.
- [32] Haq SM, Mohammed D, Waheed M, Manoj K, Manzer SH, Bussmann RW (2024) Predicting potential invasion risks of *Leucaena leucocephala* (Lam.) de Wit in the arid area of Saudi Arabia. *J. Arid Land.*, 16: 983–999. <https://doi.org/10.1007/s40333-024-0020-4>
- [33] Halmy M.W, Fawzy M, Ahmed DA, Saeed NM, Awad MA (2019) Monitoring and predicting the potential distribution of alien plant species in arid ecosystem using remotely-sensed data. *Remote Sens. Applications: Society Envi.*, 13: 69-84. <https://doi.org/10.1016/j.rsase.2018.10.005>
- [34] Hammad S, Mohamed A, Kamal M (2024) Ecophysiological adaptation and potential of energy production of two halophytes grown in the Red Sea coast of Egypt. *Egypt. J. Bot.*, 64: 189-199. DOI:10.21608/ejbo.2024.279525.2781
- [35] Hegazy AK, Hammouda O, Lovett-Doust J, Gomaa NH (2008) Population dynamics of *Moringa peregrina* along altitudinal gradient in the north- western sector of the Red Sea. *J. Arid Envi.*, 72: 1537-1551. <https://doi.org/10.1016/j.jaridenv.2008.03.001>
- [36] Hegazy A K, Fahmy AG, Mohamed HA (2004a) Shayeb El-Banat Mountain group of the Red Sea coast: a proposed biosphere reserve. *Proceedings of the symposium on natural resources and their conservation in Egypt and Africa.*, 19-21 March 2001, Cairo Univ. pp. 94-121.
- [37] Hegazy AK, Lovett-Doust J, Hammouda O, Gomaa NH (2007) Vegetation distribution along the altitudinal gradient in the northwestern Red Sea region. *J. Community Ecol*, 8 (2): 151-162. DOI:10.1556/ComEc.8.2007.2.2
- [38] Hegazy AK, Alatar AA, Lovett-Doust J, El-Adawy HA (2012) Spatial and temporal plant phenological Niche differentiation in the Wadi Degla Desert Ecosystem (Egypt). *Acta Bota. Croatica*, 71: 261-77. DOI:10.2478/v10184-011-0057-2
- [39] Hussein EA, Abd El-Ghani MM, Hamdy RS, Shalabi LF (2021) Do Anthropogenic Activities Affect Floristic Diversity and Vegetation Structure More Than Natural Soil Properties in Hyper-Arid Desert Environments?. *Diversity*, 13: 157. <https://doi.org/10.3390/d13040157>

- [40] Jackson ML (1962) Soil chemical analysis constable. Constable and Co. Ltd., London.
- [41] Jarocinska AM, Kacprzyk M, Marcinkowska-Ochtyra A, Ochtyra A, Zagajewski B, Meuleman K (2016) The application of APEX images in the assessment of the state of non- forest vegetation in the Karkonosze Mountains. *Miscellanea Geographica – Regional Studies on Development*, 20 (1): 21–27. DOI:10.1515/mgrsd-2016-0009
- [42] Kassas M (1952) Habitat and plant communities in the Egyptian Desert. I-Introduction. *J. Ecol.*, 40: 342-351. <https://doi.org/10.2307/2256804>
- [43] Kassas M (1966) Plant life in deserts. In *Arid Lands*. (E. S. Hills, ed.), pp. 145–80. UNESCO, London, Mathuen/Paris. 145-180. eBook ISBN 9781003426639
- [44] Kassas M (1971) The plant world. Part 2. Plant life in the Eastern Sahara. The Sahara and its outskirts (*I. Physiogeographie*. (H. Sciffers, ed.), pp. 477–497. Afrika-Studien, München: Weltforum-Verlag.
- [45] Khdery G A, Arafat S, Aboelghar M, Tantawy M, Abou-El-Enain M, Kamal U (2017) Selection of Optimal Hyperspectral Vegetation Indices For Estimating Chlorophyll Content of Some Plant Species. *European Chemical Bulletin*, 6:531-535. DOI:10.17628/ecb.2017.6.531-535
- [46] Khdery G, Frag E, Arafat S (2019) Natural Vegetation Cover Discrimination Using Hyperspectral Data in Wadi Hagul, Egypt. *Egypt. J. Remote Sens. Space Sci.* <https://doi.org/10.1016/j.ejrs.2019.09.002> 3
- [47] Khdery G A, Helal b, Saleh HA (2023) Using hyperspectral indices for investigation polluted plants and soil in Abu Zabaal area, Egypt. *Acta Ecologica Sinica*, 43:776-784.
- [48] Krumbein WC, Sloss LL (1963) Stratigraphy and sedimentation. Second edition. Freeman. San Francisco, pp. 497.
- [49] Kupková L, Červená L, Suchá R, Jakešová L (2017) Classification of Tundra Vegetation in the Krkonoše Mts. National Park Using APEX, AISA Dual and Sentinel-2A Data. *Eur. J. Remote Sens.*, 50: 29–46. DOI:10.1080/22797254.2017.1274573
- [50] Kurd MA, El-Khouly AA, Shawky RA, Hamad-Allah NM (2017) Vegetation-soil relationship associated with *Moringa peregrina* (Forssk.) Fiori. Population in Egypt. *Mansoura J. of Envi. Sci.*, 46: 121–136.
- [51] Kycko M, Zagajewski B, Kozłowska A (2014) Variability in spectral characteristics of trampled high-mountain grasslands. *Miscellanea Geographica – Regional Studies on Development*, 18 (2): 10–14.
- [52] Mahklouf MH (2019) Invasive Alien Plant Species In Libya. *J. of Advanced Botany and Zoology*, 7 (1): 1-4. V7I106. DOI:10.5281/zenodo.2575522. <http://scienceq.org/j.s/JABZ.php>
- [53] Mascarini L, Lorenzo G, Vilella F (2006) Leaf Area Index, Water Index, and Red: Far Red Ratio Calculated by Spectral Reflectance and its Relation to Plant Architecture and Cut Rose Production. *J. Amer. Soc. Hort. Sci.*, 131: 313–319. DOI:10.21273/JASHS.131.3.313
- [54] Nagendra H (2002) Using remote sensing to assess biodiversity. *Int. J. Remote Sens.*, 22(12): 2377–2400. DOI:10.1080/01431160117096
- [55] Raunkiaer C (1934) The life forms of Plants and statistical plant geography, being the collected papers of C. Raunkiaer. Clarendon Press, Oxford, GB, 632 pp.
- [56] Richards LA (1954). Diagnosis and improvement of saline and alkali soils. *Soil Sci.*, 78 (2): 1-154.
- [57] Ryan J, Garabet S, Harmson K, Rashid A (1996) A soil and plant analysis. Manual adapted for the West Asia and North Africa region. ICARDA Tech. Bull., pp. 244.
- [58] Salama FM, Abd El-Ghani MM, Amro A, Gaafar A, Abd El Galil AA (2018) Vegetation Dynamics and Species Diversity in a Saharan Oasis, Egypt. *Notulae Scientia Biologicae*, 10: 363-372. DOI:10.15835/nsb10310296
- [59] Salama FM, Abd El-Ghani MM, El-Tayeh NA, Amro AM, Gaafar AA, Abd El- Galil AA (2019) Assessing the role of environmental gradients on the phytodiversity in Kharga Oasis of Western Desert, Egypt. *Jordan J. Biological Sci.*, 12: 421-434.
- [60] Salama FM, Abd El-Ghani M, Gadallah M, El-Naggar S, Amro A (2014) Variations in Vegetation Structure, Species Dominance and Plant Communities in South of the Eastern Desert-Egypt. *Notulae Scientia Biologicae*, 6 (1): 41-58. DOI:10.15835/nsb.6.1.9191
- [61] Salama F, El-Tayeh NA, Zaher AM, El- Naggar S, Gaafar A (2023) Phytosociological Studies on the Associated Species of *Balanites aegyptiaca* in the Eastern and Western Egyptian Deserts. *Egyp. J. of Botany*, 63: 1005-1029. DOI:10.21608/ejbo.2023.205410.2308
- [62] Salama FM, Abd El-Ghani MM, El-Tayeh NA (2013) Vegetation and soil relationships in the inland wadi ecosystem of central Eastern Desert, Egypt. *Turk. J. Bot.* 37 (3), 489-498.
- [63] El-Khouly AA, Shawky R (2017) Plant species diversity of some wadis at Red Sea Coast, Egypt. *Research & Reviews: J. of Botanical Sciences*, 6: 25-35. DOI:10.23977/erej.2017.11001
- [64] Shaltout KH, Sheded MG, Salem AH (2009) Population structure of common shrubs and trees in Wadi Allaqi Biosphere Reserve, South-East Egypt. *Feddes Report* 120: 343–354. DOI:10.1002/fedr.200911114
- [65] Sheded MG, Hamed ST, Badry MO (2014) Vegetation analysis six riverian islands in hyper-Arid environments

at Qena Government (Upper-Egypt). *Acta Botanica Hung.*, 56 (304): 409-431. DOI:[10.1556/ABot.56.2014.3.4.15](https://doi.org/10.1556/ABot.56.2014.3.4.15)
 [66] Täckholm V (1974) Students' Flora of Egypt, 2nd edition. Cairo University Publisher, Cooperative Printing Company, Beirut, 888 p.

[67] Wisz MS, Pottier J, Kissling WD, Pellissier L, Lenoir J, Damgaard CF, Wardle D, Aastrup P, Svenning J C et al. (2013) The role of biotic interactions in shaping distributions and realized assemblages of species: implications for species distribution modeling. *Biol. Rev. of the Cambridge Philosophical Society*, 88:15–30. Doi: [10.1111/j.1469-185X.2012.00235.x](https://doi.org/10.1111/j.1469-185X.2012.00235.x)

[68] Zaghloul MS, Abd El-Wahab RH, Moustafa AA (2010) Ecological assessment and phenotypic and fitness variation of Sinai's remnant populations of *Moringa Peregrina*. *Appl. Ecol. Environ. Res.*, 8 (4): 351–366. <http://www.ecology.uni-corvinus.hu>.

[69] Zahran MA (1982) Ecology of the halophytic vegetation of Egypt. In: Sen, D.N., Rajpurohit, K.S. (eds) Contributions to the ecology of halophytes. Tasks for vegetation science, vol 2. Springer, Dordrecht. https://doi.org/10.1007/978-94-009-8037-2_2

[70] Zahran MA, El-Amier YA (2014) Ecology and establishment of fiber producing taxa naturally growing in the Egyptian deserts. *Egypt. J. Basic Appl. Sci.*, 1:144–150. DOI:[10.1016/j.ejbas.2014.10.001](https://doi.org/10.1016/j.ejbas.2014.10.001)

[71] Zahran MA, Willis AJ (2009) The Vegetation of Egypt, Plant and Vegetation. Springer. <https://doi.org/10.1007/978-1-4020-8756-1>.




Fig. 1. Location map of the study area showing the studied three mountains (Gebel El-Sibaie, Gebel Umm lasifa and Gebel Umm Naqqat) and their main Wadis (Esel, Sharm El-Bahari and Umm Gheig)




Fig. 2. Life span spectrum of the total study area vegetation




Fig. 3. Histogram of the floristic composition of the 29 different families surveyed in Gabal El-Sebaie area.




Fig. 4. Life form spectrum of the of the 60 plant species recording in the total study area, showing five different life forms.




Fig. 5. Life form histogram and percentage of species at 6 study area locations: Gebel El-Sibaie (Se), Gebel Umm lasifa (La), Gebel Umm Naqqat (Na), Wadi Esel (Es), Sharm El-Bahari (Sh) and Umm Gheig(Gh).





Fig. 6. Phytochorion categories of the total recorded species and their relevant percent (%) in Jabal El-Sebaie area. (Am= American, AUS=Australian, SA-AR= Saharo- Arabian, SU-ZA= Sudano-Zambezian, ME= Mediterranean, IR-TR= Irano-Turanian, Pan = Pantropical, AF-MO = Afro-Montane, SU=Sudanien, ER-SR = Euro-Siberian and SA-SI= Saharo-Sindian). and percentage of species.






Fig.8. ANOVA and Tukey's HSD analyses used to differentiate the studied taxa




Fig.9. Spectral reflectance patterns of the top soil in the examined soil profiles.




Fig.10. ANOVA and Tukey's HSD analysis of soil types across various spectral bands (Blue, Green, NIR and Red Zones, SWIRI and SWIRII).

Table 1. The number of recorded species belonging to the main floristic chorotypes and their r and their relevant percent (%) in Jabal El-Sebaie area. Phytochoria categories (Am= American, AUS=Australian, SA-AR= Saharo- Arabian, SU-ZA= Sudano-Zambezian, ME= Mediterranean, IR-TR= Irano-Turanian, Pan = Pantropical, AF-MO = Afro-Montane, SU=Sudanian, ER-SR = Euro-Siberian and SA-SI= Saharo-Sindian).

Phytocchoria categories	No. of plant species	Percentage (%)
Monoregional		
AUS	1	1.67
AM	2	3.33
SU	1	1.67
ME	1	1.67
SA-AR	15	25.00
SU-ZA	3	5.00
SA-SI	6	10.00
Total	29	48.33
Bioregional		
ME + SA-AR	1	1.67
SA-AR + SU	1	1.67
SA-AR + SU-ZA	13	21.67
SA-AR +IR-TR	7	11.67
Total	22	36.67
Pluriregional		
ME+SA-AR+IR-TR	2	3.33
ME+SA-AR+SU-ZA	1	1.67
ME+IR-TR+ER-SR	1	1.67
ME+SA-SI+IR-TR	1	1.67
SA-AR+SU-ZA+IR-TR	1	1.67
Total	6	10.00
Others		
Pan	2	3.33
AF-MO	1	1.67
Total	3	5.00

Appendix1: Species composition of the study area classified according to the different families, together with their chorology types, and the distribution of species in the studied Area. Gebel El-Sibaie (Se), Gebel Umm-Naqqat (Na), Gebel Umm-Lasifa (La), Wadi Sharm El-Bahari (Sh), Wadi Umm-Gheig (Gh), and Wadi Esel (Es). Life forms (Th: Therophytes, He: Hemicryptophytes, Ch: Chamaephytes, G: Geophytes and Ph: Phanerophytes)

	Family	Species incl. author	Life cycle	Life form	Growth form	Chorology	Study area					
							Se	Na	Gh	La	Sh	Es
1	Aizoaceae	<i>Aizoon canariense</i> L.	A	Th	herb	SU-ZA		*	*			
2	Amaranthaceae	<i>Aerva javanica</i> (Burm. f.) Juss. ex Schult. in Roem. & Schult. Var <i>javanica</i>	P	Ch	herb	SA-AR+S U-ZA		*				

4	Asclepiada ceae	<i>Leptadenia pyrotechnica</i> (Forss k.) Decne.	P	Ph	shru b	SA- AR+S U-ZA	*	*				*
5		<i>Solenostemma arghel</i> (Delile) Hayne	P	Ch	shru b	SA-AR	*	*	*			
6	Asteraceae	<i>Artemisia judaica</i> subsp. <i>juda ica</i>	P	Ch	shru b	SA-AR		*				
7		<i>Launaea capitata</i> (Spreng.) Dandy	A	Th	Her b	SA-AR			*			
8		<i>Pulicaria undulata</i> (L.) C.A.Mey.	P	C h	shru b	SA- AR+S U-ZA		*				
9		<i>Pulicaria incisa</i> (Lam.) DC.	P	C h	herb	SA- AR+I R-TR	*					
1 0		<i>Senecio flavus</i> (Decne.) Sch.Bip.	A	Th	herb	SA-SI	*	*				
1 1	Avicennia ceae	<i>Avicennia marina</i> (Forssk.) Vierh.	P	Ph	tree	SA-AR					*	
1 2	Boraginac eae	<i>Arnebia hispidissima</i> (Siebe r ex Lehm.) A.DC.	A	Th	herb	SA- AR+I R-TR	*		*			
1 3		<i>Trichodesma ehrenbergii</i> Schwei nf.	P	H	herb	SU-ZA	*	*				
1 4	Brassicace ae	<i>Zilla spinosa</i> (L.) subsp. <i>spinosa</i>	P	Ch	shru b	SA-AR	*	*	*			*
1 5	Capparace ae	<i>Capparis decidua</i> (Forssk.) Edgew.	P	Ph	tree	SA- AR+S U	*		*			*
1 6		<i>Capparis sinaica</i> Veill	P	Ch	shru b	SA-SI	*	*	*	*		*
1 7	Caryophyl laceae	<i>Paronychia argentea</i> Lam.	A	Th	herb	ME			*			
1 8		<i>Cometes abyssinica</i> R.Br. in Wall.	A	Th	herb	AF- MO		*	*			*
		<i>Polycarpaea repens</i> (Forssk.) Asch. & Schweinf.	P	H	herb	SA-SI			*			

1 9		<i>Spergularia diandra</i> (Guss.) Boiss .	A	Th	herb	ME+S A- SI+IR- TR			*			
	Chenopod iaceae	<i>Salsola imbricata</i> Forssk.	P	Ch	shru b	SA-AR			*		*	*
2 0	Cleomace ae	<i>Cleome amblyocarpa</i> Barrat te & Murb.	A	Th	herb	SA- AR+S U-ZA			*			*
2 1		<i>Cleome chrysantha</i> Decne.	p	Ch	shru b	SA-SI	*					
2 2		<i>Cleome droserifolia</i> (Forssk.) Delile	P	Ch	shru b	SA- AR+I R-TR	*	*	*			*
2 3	Convolvul aceae	<i>Convolvulus hystrix</i> Vahl subsp. <i>hystrix</i>	P	C h	shru b	SA-AR		*	*			*
2 4	Cucurbita ceae	<i>Citrullus colocynthis</i> (L.) Vahl .	P	H	herb	ME+S A-AR	*	*				
2 5	Fabaceae	<i>Acacia cyclocarpa</i> Maslin, M.D.Barrett & R.L.Barrett	P	Ph	tree	AUS						*
2 6		<i>Acacia ehrenbergiana</i> Hayne.	P	Ph	tree	SU						*
2 7		<i>Acacia tortilis</i> (Forssk.) Hayne subsp. <i>raddiana</i> (Savi) Brenan	P	Ph	tree	SA- AR+S U-ZA	*	*	*	*		*
2 8		<i>Acacia tortilis</i> (Forssk.) Hayne subspecies <i>tortilis</i> (Forssk.)	P	Ph	tree	SA- AR+S U-ZA	*	*	*			*
2 9		<i>Astragalus eremophilus</i> Boiss.	A	Th	herb	SA-AR			*			
3 0		<i>Astragalus vogelii</i> (Webb) Bornm.	A	Th	herb	SA-AR			*			
3 1		<i>Leucaena leucocephala</i> (Lam.) de Wit	P	Ph	shru b	AM						*
3 2		<i>Lotus glinoides</i> Delile	A	Th	herb	SA-SI			*			
3 3		<i>Prosopis juliflora</i>	P	Ph	shru b	AM						*

3 4		<i>Taverniera aegyptiaca</i> Boiss.	P	C h	shru b	SA-AR			*		*	*
3 5		<i>Tephrosia purpurea</i> (L.) Pers. subsp. <i>apolli</i> nea (Delile) Hosni & El-Karemy	P	C h	herb	SA-AR	*	*				
3 6	Liliaceae	<i>Asphodelus tenuifolius</i> Cav.	A	Th	herb	ME+S A- AR+I R-TR		*				
3 7	Malvaceae	<i>Malva parviflora</i> L.	A	Th	herb	ME+I R- TR+E R-SR						*
3 8	Moringaceae	<i>Moringa peregrina</i> (Forssk.) Fiori	P	Ph	tree	SU-ZA	*	*		*		
3 9	Nitrariaceae	<i>Nitraria retusa</i> (Forssk.) Asch.	P	Ph	shru b	SA- AR+S U-ZA		*	*			*
4 0	Plumbaginaceae	<i>Limonium axillare</i> (Forssk.) Kuntze	P	C h	shru b	SA- AR+S U-ZA					*	
4 1	Poaceae	<i>Aristida funiculata</i> Trin. & Rupr.	A	Th	grass	SA-AR			*			
4 2		<i>Panicum turgidum</i> Forssk.	P	G	grass	SA- AR+I R-TR		*	*			
4 3		<i>Phragmites australis</i> (Cav.) Trin. ex Steud.	P	G	shru b	Pan					*	
4 4	Resedaceae	<i>Ochradenus baccatus</i> Delile	P	Ph	shru b	SA- AR+S U-ZA	*	*	*			*
4 5		<i>Reseda pruinosa</i> Delile	A	Th	herb	SA-AR	*	*	*			
4 6	Rhamnaceae	<i>Ziziphus spina-christi</i> (L.) Desf.	P	Ph	tree	SA- AR+S U-ZA	*					
4 7	Rubiaceae	<i>Crucianella membranacea</i> Boiss.	A	Th	herb	SA-AR						*

48	Salvadoraceae	<i>Salvadora persica</i> L.	P	Ph	shrub	SA-AR+S U-ZA		*	*			
49	Scrophulariaceae	<i>Kickxia acerbiana</i> (Boiss.) Tackh.&Boulos.	P	He	herb	SA-SI	*					
50	Solanaceae	<i>Lycium shawii</i> Roem. & Schult.	P	Ph	shrub	SA-AR+S U-ZA+IR-TR	*	*	*			*
51	Tamaricaceae	<i>Tamarix nilotica</i> (Ehrenb.) Bunge	P	Ph	tree	SA-AR+I R-TR					*	*
52	Urticaceae	<i>Forsskaolea tenacissima</i> L.	P	H	herb	SA-AR+S U-ZA	*					
53	Zygophyllaceae	<i>Balanites aegyptiaca</i> (L.) Delile	P	Ph	tree	SA-AR+S U-ZA	*	*	*			*
54		<i>Fagonia arabica</i> L. var. <i>arabica</i>	P	Ch	herb	SA-AR+I R-TR	*	*				
55		<i>Fagonia mollis</i> Delile	P	Ch	shrub	SA-AR+I R-TR	*	*	*			
56		<i>Tribulus terrestris</i> L.	A	Th	herb	Pan		*				*
57		<i>Zygophyllum album</i> L.f.	P	Ch	herb	ME+S A-AR+I R-TR			*		*	*
58		<i>Zygophyllum coccineum</i> L.	P	Ch	shrub	ME+S A-AR+S U-ZA	*		*		*	*
59		<i>Zygophyllum simplex</i> L.	A	Th	herb	SA-AR	*		*			

Appendix (2). Means and standard deviations for soil physical properties of the twenty-four sites at depth (0-20cm) in the study area. ANOVA test is for original where values are normally distributed. F-value refer to the ANOVA. Small letters denote the statistically different sites as identified by ANOVA

Sites	Habitat	Particle size distribution (%)					
		Very coarse sand	Coarse sand	Medium sand	Fine sand	Very fine Sand	Silt & Clay

El-Sebaie mountain	Slope	^d 46.23 ± 3.0820	^{bij} 18.15 ± 2.0167	^f 14.03 ± 1.5589	^h 9.08 ± 1.0089	^{gh} 6.48 ± 0.7200	^{gh} 6.03 ± 0.6700
		^{jk} 19.36 ± 1.2907	^{gh} 21.64 ± 2.4044	^e 17.64 ± 1.9600	^e 12.76 ± 1.4178	^{ef} 16.38 ± 1.8200	^d 12.22 ± 1.3578
		^{hi} 26.57 ± 1.7713	^a 47.49 ± 5.2767	^{de} 20.06 ± 2.2289	^l 3.62 ± 0.4022	ⁱ 1.53 ± 0.1700	^{jk} 0.73 ± 0.0811
		^b 59.63 ± 3.9753	^{de} 29.56 ± 3.2844	^{hi} 8.85 ± 0.9833	^{ijk} 1.25 ± 0.1389	^l 0.44 ± 0.0489	^k 0.27 ± 0.0300
		^{ef} 37.89 ± 2.5260	^{ijk} 15.92 ± 1.7689	^a 27.64 ± 3.0711	^h 7.67 ± 0.8522	^{gh} 6.00 ± 0.6667	^{hi} 4.88 ± 0.5422
		^b 61.03 ± 4.0687	^{cde} 30.84 ± 3.4267	^l 6.64 ± 0.7378	^{jk} 0.84 ± 0.0933	^l 0.35 ± 0.0389	^k 0.30 ± 0.0333
	Wadi bed	^o 2.62 ± 0.1747	^l 3.79 ± 0.4211	^l 6.18 ± 0.6867	^d 15.97 ± 1.7744	^b 38.58 ± 4.2867	^e 32.86 ± 3.6511
		^c 53.32 ± 3.5547	^{bc} 36.04 ± 4.0044	^l 6.80 ± 0.7556	^{ijk} 1.46 ± 0.1622	^l 1.95 ± 0.2167	^k 0.43 ± 0.0478
		^{mn} 10.00 ± 0.6667	^k 10.58 ± 1.1756	^e 17.34 ± 1.9267	^e 20.02 ± 2.2244	^e 32.03 ± 3.5589	^{def} 10.03 ± 1.1144
		^{fg} 33.52 ± 2.2347	^b 39.23 ± 4.3589	^{cde} 20.29 ± 2.2544	^{ij} 3.22 ± 0.3578	^l 2.04 ± 0.2267	^{jk} 1.70 ± 0.1889
		^{hi} 26.64 ± 1.7760	^{fgh} 22.13 ± 2.4589	^{gh} 10.82 ± 1.2022	^{fgh} 10.10 ± 1.1222	^{ef} 19.30 ± 2.1444	^{de} 11.01 ± 1.2233
		^o 1.45 ± 0.1611	^l 1.63 ± 0.1811	^j 1.67 ± 0.1856	^{efg} 1.71 ± 1.3011	^a 45.85 ± 5.0944	^b 37.69 ± 4.1878
Wadi Esel	Wadi bed	ⁿ 7.63 ± 5.8692	^{jk} 14.71 ± 1.6344	^{ab} 25.84 ± 2.8711	^a 24.43 ± 2.7144	^e 19.75 ± 2.1944	^{fg} 7.64 ± 0.8489
Umm Lasifa mountaion	Slope	^b 64.39 ± 4.2927	^{efg} 26.31 ± 2.9233	^l 6.06 ± 0.6733	^{ijk} 1.55 ± 0.1722	^l 0.38 ± 0.0422	^{jk} 1.31 ± 0.1456
Wadi Sharm El-Bahary	Mangrove	^o 2.77 ± 0.1847	^k 10.48 ± 1.2200	^{hi} 7.83 ± 0.8700	^{ef} 11.93 ± 1.3256	^d 25.05 ± 2.7833	^a 41.44 ± 4.6044
	Wadi bed	^{mn} 12.06 ± 0.8040	^{bij} 17.18 ± 1.9089	^{bcd} 22.72 ± 2.5244	^{bc} 21.51 ± 2.3900	^{rf} 17.82 ± 1.9800	^{efg} 8.71 ± 0.9678
Umm Naqqat mountain	Slope	^{de} 42.28 ± 2.8187	^{bc} 36.08 ± 4.0089	^{fg} 12.11 ± 1.3456	^l 3.63 ± 0.4033	^{hi} 3.46 ± 0.3844	^{ijk} 2.44 ± 0.2711
		^a 75.75 ± 5.0500	^{hi} 20.34 ± 2.2600	ⁱ 2.54 ± 0.2822	^k 0.39 ± 0.0433	^l 0.47 ± 0.0522	^{jk} 0.51 ± 0.0567
		^{ij} 23.5 ± 1.5667	^{def} 27.13 ± 3.0144	^b 24.11 ± 2.6789	^{ef} 12.52 ± 1.3911	^g 9.26 ± 1.0289	^{bij} 3.48 ± 0.3867
	Wadi bed	^{de} 42.12 ± 2.8080	^a 48.85 ± 5.4279	^{hi} 8.51 ± 0.9454	^k 0.43 ± 0.0478	^l 0.05 ± 0.0056	^k 0.04 ± 0.0044
		^d 47.04 ± 3.1360	^{cde} 31.48 ± 3.4978	^f 14.21 ± 1.5789	^{ij} 3.27 ± 0.3633	^l 2.31 ± 0.2567	^{jk} 1.69 ± 0.1878
		^{gh} 29.22 ± 1.9480	^{cd} 32.02 ± 3.5578	^b 24.22 ± 2.6911	^{gh} 9.40 ± 1.0444	^{hi} 3.12 ± 0.3467	^{jk} 2.02 ± 0.2244
Wadi Umm Gheig	Wadi bed	^{kl} 14.96 ± 0.9973	^{jk} 13.33 ± 1.4811	^{bc} 23.43 ± 2.6033	^{ab} 22.99 ± 2.5544	^f 16.18 ± 1.7978	^{ef} 9.11 ± 1.0122

		^{lm} 12.96 ± 0.8640	^{bij} 17.69 ± 1.9656	^{bc} 23.51 ± 2.6122	^{±bc} 21.88 ± 2.4311	^{±f} 16.12 ± 1.7911	^{±fg} 7.84 ± 0.8711
F- value		180.1803	52.483	56.792	100.686	136.348	164.561
LSD _{0.05}		4.582	4.891	3.046	2.267	3.185	14.997

Appendix (3). Means and standard deviations for soil chemical properties of the twenty-four sites at depth (0-20cm) in the study area. ANOVA test is for original where values are normally distributed. F-value refer to the ANOVA. Small letters denote the statistically different sites as identified by ANOVA

Location	Habitat	EC ds/m	PH	TDS Mg / L	Soluble cations (meq / L)				Soluble anions (meq / L)			P mg / L
					Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	HCO ₃ ⁻	SO ₄ ⁻²	Cl ⁻	
El-Sebaie mountain	Slope	^{ij} 0.28 ^b ± 0.0075	8.79 ± 0.0068	^r 175.61 ± 0.3856	^g 1.53 ± 0.0411	^g 0.20 ± 0.0055	^{mno} 0.64 ± 0.0172	^{±k} 0.43 ± 0.0115	^{±fg} 1.23 ± 0.0433	^{±l} 1.02 ± 0.0358	^{±j} 0.58 ± 0.0204	^{gh} 18.09 ± 0.3547
		^{hi} 0.71 ^k ± 0.0231	8.00 ± 0.0033	ⁿ 430.66 ± 0.4929	^f 4.89 ± 0.1580	^g 0.82 ± 0.0264	^{lm} 0.94 ± 0.0303	^{hi} 0.50 ± 0.0162	^{efg} 2.59 ± 0.1180	^{±l} 2.13 ± 0.0971	^h 2.54 ± 0.1157	^o 5.59 ± 0.1098
		^j 0.22 ^e ± 0.0182	8.50 ± 0.0045	^w 125.67 ± 0.3627	^g 0.79 ± 0.0662	^g 0.30 ± 0.0249	^{lmn} 0.91 ± 0.0765	^{mn} 0.17 ± 0.0143	^{±fg} 1.22 ± 0.0731	^{±l} 0.40 ± 0.0241	^{±j} 0.49 ± 0.0292	^{jk} 14.19 ± 0.2782
		^j 0.23 ^h ± 0.0056	8.20 ± 0.0034	^v 129.90 ± 0.4567	^g 1.02 ± 0.0253	^g 0.54 ± 0.0134	^o 0.51 ± 0.0127	^{mn} 0.18 ± 0.0043	^{±fg} 1.23 ± 0.0520	^{±l} 0.47 ± 0.0197	^{±j} 0.61 ± 0.0256	^k 13.67 ± 0.2680
		^{ij} 0.35 ^e ± 0.0036	8.50 ± 0.0031	^p 224.43 ± 0.3286	^g 2.10 ± 0.0211	^g 0.11 ± 0.0011	^{lmno} 0.88 ± 0.0088	^{±j} 0.46 ± 0.0047	^{±g} 1.12 ± 0.0021	^{±l} 1.75 ± 0.0033	^{±j} 0.71 ± 0.0014	^j 14.55 ± 0.2853
		^{ij} 0.36 ⁿ ± 0.0023	7.50 ± 0.0049	^q 215.90 ± 0.7962	^g 1.99 ± 0.0127	^g 0.42 ± 0.0027	^{kl} 1.05 ± 0.0067	^{mn} 0.18 ± 0.0011	^{±fg} 1.40 ± 0.0063	^{±l} 1.33 ± 0.0060	^{±ij} 0.95 ± 0.0043	^r 0.0049 ± 0.001

	Wadi bed	^g 1.41 ^j 8.10 ± 0.060 8	^j 846.09 ± 6.55 0.9253 0.2831	^g 1.31 ± 4.12 ^b 2.09 0.0567 0.177 0.090	ⁱ 4.12 ^b 2.09 0.0567 0.177 0.090	^{def} 3.31 ± 7.96 ^h 2.67 ^e 22.2 0.2000 0.480 0.161 0.43 8 5 65
		^{gh} 1.07 ± 0.0046 ⁿ 7.49 0.005 0.4072	^f 661.92 ± 5.80 0.0248 0.0053	^j 1.95 ^c 1.68 0.008 0.007	^{de} 3.77 ± 4.56 ^k 2.36 ^c 30.0 0.0211 0.025 0.013 0.58 6 2 94	
		^e 3.64 ^s 6.89 ± 0.0424 0.004 ± 0.8898	^d 2130.88 ± 18.56 0.2162 0.1323	^g 6.05 ^{jk} 0.43 0.070 0.005	^b 15.15 ± 15.55 ^f 6.41 ^a 40.2 0.438 0.180 0.78 9 9 92	
		^g 1.38 ^m 7.79 ± 0.0437 0.01 ± 0.64	^f 868.44 ± 5.86 0.6732 0.1853	^h 4.87 ^d 0.91 0.154 0.028	^{efg} 2.67 ± 7.30 0.0115 ± 0.03 16 5	^g 4.28 ^f 21.2 0.018 0.41 5 59
		^{gh} 1.01 ± 0.0113 ^p 7.19 0.005 ± 0.2536	^m 632.12 ± 5.47 0.612 0.0222	^j 2.16 ^h 0.51 0.024 0.005	^{efg} 2.36 ± 5.93 ^{jk} 1.88 ^k 13.6 0.0360 0.090 0.028 0.26 6 7 80v	
Wadi Esel	Wadi bed	^d 5.86 ^u 6.49 ± 0.0559 0.023 ± 1.2263	^e 3580.61 ± 25.81 0.2463 0.0986	^b 10.33 ± 20.76 0.198 0.015	^c 1.65 ^e 7.66 ± 26.75 0.0741 ± 0.25 7 89	^e 24.15 ± 0.65 0.23 ± 0.0 36 128
		^g 1.32 ^l 7.90 ± 0.0117 0.000 ± 0.2626	^k 823.56 ± 5.99 0.0532 0.0165	^h 4.97 ⁱ 0.38 0.044 0.003	^{fg} 1.67 ± 6.89 ^{ij} 4.65 ^r 0.00 0.0157 0.064 0.043 ± 0.0 8 8 009	
Umm Lasifa mountain	Slope	^{ij} 0.24 ^d 8.60 ± 0.0169 0.015 ± 0.4434	^u 135.96 ± 0.59 0.0417 0.0786	^{no} 0.54 ± 0.147 0.038 ± 0.010	^l 0.58 ^j 0.51 ⁿ 6.64 0.045 0.040 ± 0.13 9 6 02	
Wadi Shar m El-Bahary	Mangrove	^a 25.68 ± 1.1994 ^a 9.00 0.001 ± 7.4993	^a 14277.37 ± 96.68 4.5165 146.55 ± 6.8461	^d 12.70 ± 0.5934 ^e 0.82 0.038	^a 103.42 ± 5.1239 4.788 2.852 0.36 4 5 37	

	Wadi bed	^b 9.50 ^r ± 0.028 6	^r 7.05 ± 0.040 7	^b 6031.89 ^b ± 2.6191	^b 37.30 ^c ± 0.1124	^c 33.20 ^a ± 0.1000	^a 23.87 ^f ± 0.0719	^f 0.66 ^c ± 0.0020	^c 8.61 ^b ± 0.1198	^b 71.19 ^d ± 0.9906	^d 16.45 ^b ± 0.2289	^b 34.2 ³ ± 0.6712
Umm Naqqat mountain	Slope	^{ij} 0.40 ^l ± 0.011 0	^l 8.16 ^s ± 0.039 7	^s 251.00 ^{±g} ± 0.2569	^{±g} 1.98 ^{±g} ± 0.0544	^{±g} 0.29 ^{±±} ± 0.0081	^k 1.35 ^{kl} ± 0.0370	^{kl} 0.40 ^{fg} ± 0.0108	^l 1.26 ^{±±} ± 0.0035	^l 2.10 ^j ± 0.0059	^j 0.71 ^m ± 0.0020	^m 7.42 ¹ ± 0.1455
		^{ij} 0.28 ^c ± 0.003 6	^c 8.65 ^s ± 0.050 3	^s 168.60 ^{±g} ± 0.3090	^{±g} 1.44 ^{±g} ± 0.0185	^{±g} 0.60 ^{±±} ± 0.0077	^{mno} 0.56 ^{mn} ± 0.0072	^{mn} 0.18 ^{±g} ± 0.0023	^l 1.10 ^{±±} ± 0.0415	^l 1.07 ^j ± 0.0407	^j 0.70 ^d ± 0.0263	^d 26.1 ⁶ ± 0.5129
		^f 3.07 ^v ± 0.127 0	^v 6.26 ^g ± 0.052 4	^g 1773.17 ^e ± 1.7242	^e 14.10 ^c ± 0.5836	^c 10.09 ^{±±} ± 0.4177	^g 5.95 ^{±±} ± 0.2461	^g 0.55 ^{±±} ± 0.0227	^b 14.60 ^{±±} ± 0.9003	^h 8.86 ^e ± 0.5463	^e 8.01 ^h ± 0.4938	^h 17.8 ³¹ ± 0.3496
	Wadi bed	^{ij} 0.26 ^g ± 0.007 3	^g 8.40 ^r ± 0.002 9	^r 149.67 ^{±g} ± 0.5997	^{±g} 0.91 ^{±g} ± 0.0257	^{±g} 0.81 ^{±±} ± 0.0229	^{lmno} 0.68 ^m ± 0.0193	^m 0.20 ^{fg} ± 0.0057	^{fg} 1.23 ^{±±} ± 0.0442	^l 0.64 ^j ± 0.0229	^j 0.75 ^p ± 0.0270	^p 3.77 ⁷ ± 0.0741
		^{ij} 0.25 ^f ± 0.009 6	^f 8.45 ^r ± 0.045 7	^r 148.36 ^{±g} ± 0.8476	^{±g} 1.30 ^{±g} ± 0.0493	^{±g} 0.22 ^{±±} ± 0.0082	^{lmno} 0.82 ^{mn} ± 0.0310	^{mn} 0.19 ^{±g} ± 0.0071	^g 1.00 ^{±±} ± 0.0163	^l 0.95 ^j ± 0.0156	^j 0.60 ^l ± 0.0099	^l 11.3 ² ± 0.2220
	Wadi Umm Gheig	Wadi bed	^f 2.96 ^r ± 0.066 5	^r 6.70 ^h ± 0.003 0	^h 1758.54 ^d ± 1.1691	^d 18.11 ^f ± 0.4063	^f 4.27 ^{±±} ± 0.0957	^f 6.70 ^g ± 0.1502	^g 0.55 ^{fg} ± 0.0123	^{fg} 1.57 ^{±±} ± 0.0241	^g 12.50 ^d ± 0.1922	^d 15.68 ⁱ ± 0.2410
^c 6.29 ^q ± 0.008 4			^q 7.09 ^d ± 0.004 6	^d 3907.94 ^c ± 0.7534	^c 25.12 ^d ± 0.0335	^d 22.90 ^{±±} ± 0.0305	^e 8.71 ^a ± 0.0116	^a 6.13 ^d ± 0.082	^d 5.04 ^{±±} ± 0.0015	^d 38.30 ^c ± 0.0114	^c 19.55 ^o ± 0.0058	^o 5.66 ² ± 0.1110
^c 6.69 ^o ± 0.256 2			^o 7.27 ^c ± 0.028 5	^c 4125.10 ^e ± 3.7201	^e 13.71 ^b ± 0.5253	^b 35.98 ^{±±} ± 1.3788	^c 17.02 ⁿ ± 0.6520	ⁿ 0.15 ^d ± 0.0057	^d 4.93 ^{±±} ± 0.1917	^c 45.76 ^d ± 1.7797	^d 16.22 ^l ± 0.6307	^l 11.4 ² ± 0.2239

F- value	1391.7455	3434.9918	7785170.8	1403.9216	1360.5412	3310.3176	8918.7742	1130.438	1566.7158	1298.4577	2708.515
LSD _{0,05}	0.416	0.038	3.155	1.553	2.346	0.332	0.038	1.754	1.771	1.007	0.580