

Effectiveness Of Educational Interventions Integration With Health Services In Improving Oral Health Behaviors And Preventing Oral Diseases; Systematic Review

Noof khaled A alsahiem¹, Khlood Mohammad Alasiri², Nourah Falah Al-Mutairi³, Aljawharah Suliman Alomair⁴, Nourah Nasserallah Alomran⁵

¹Senior Specialist public health, Armed Forces Hospital, Dental department, Dhahran, Saudi Arabia

^{2,3}Dental and oral health specialists, Dental Department, King Fahad Military Medical Complex, Dhahran, Saudi Arabia

⁴Dental Hygienist, Dental Department, Armed Forces Hospital, Dhahran, Saudi Arabia

⁵Dental and oral health specialist, Dental Department, Armed Forces Hospital, Dhahran, Saudi Arabia

Abstract

Background: Oral diseases is a major global health concern, mainly in children and adolescents. Integrating educational interventions within health services can improve oral health behaviors and clinical outcomes. In this systematic review we aimed to evaluate the effectiveness of educational interventions integrated with health services to improve oral health behaviors and prevent oral diseases.

Methods: A literature search was conducted in PubMed, Scopus, Web of Science, and Google Scholar. We include studies with individual's ≤ 19 years, include educational strategies linked to health systems, and report clinical oral health outcomes. Thirteen studies were included following PRISMA guidelines. Data were synthesized qualitatively. Methodological quality was assessed using the Joanna Briggs Institute (JBI) appraisal tools.

Results: Included interventions differ in type (digital tools, school-based sessions, visual aids) and delivery settings (schools, primary care, community clinics). Most interventions show improvements in knowledge, brushing behavior, plaque index, gingival health, or ECC reduction. Integration within health services and reinforcement by health professionals or teachers improve effectiveness.

Conclusion: Educational interventions integrated into health systems promote oral health behaviors and prevent disease. Implementation in school and primary care is recommended for long-term public health improvement.

Keywords: Oral health education, Health services integration, Preventive dentistry, Behavioral intervention

INTRODUCTION

Oral diseases is a widespread global problem, contributing to pain, discomfort, and affect quality of life for millions of individuals (1). Conventional oral health education show benefits, and integrating educational interventions within health service systems also show greater potential in to promote oral hygiene practices and prevent oral conditions (2). This integration provide a systematic, accessible, and contextually relevant approach to oral health promotion, mainly in vulnerable populations (3).

Combination of oral health education with primary care centers, community clinics, and school health services, facilitates routine exposure to prevent messages and reinforces behavior through interaction with healthcare professionals (2). This aligns with the strategy of addressing common risk factors for chronic diseases, creating synergy between oral and general health initiatives (4). The agreement between oral health promotion and general health services fosters a more holistic and cost-effective model of care, mainly in settings where dental services alone are not available (4).

School-based interventions, mainly offer opportunities to educate children in their formative years while involving teachers and parents (5). These programs shape lifelong habits that prevent dental caries and periodontal conditions (5). Community-based programs embedded in health service contexts enhance oral health literacy and empower individuals to take control of their preventive care practices (3).

Strategic scaling of integrated educational interventions is important to address equity gaps and oral health outcomes on a broader scale (2). This review aims to evaluate the effectiveness of educational interventions integrated with health services to improve oral health behaviors and prevent oral diseases.

METHOD

Search Strategy

A systematic literature search was conducted to identify interventional studies which evaluate educational strategies and aimed to improve oral health in children and adolescents. Databases searched include PubMed, Scopus, Web of Science, and Google Scholar. Keywords include, oral health education, school-based intervention, children, adolescents, plaque index, and preventive dentistry. We also do manual searches of references in relevant articles to identify additional eligible studies. Only full-text, peer-reviewed articles published in English were considered.

Inclusion Criteria

We include studies when, the intervention involved an educational strategy targeting oral health behavior or awareness, participants were children or adolescents aged ≤ 19 years, measured oral health-related outcomes, design was experimental or quasi-experimental, full-text was accessible. We exclude reviews, opinion pieces, or editorials, studies included adult participants exclusively, and studies did not assess behavioral or clinical oral health outcomes.

Study Selection

Two independent reviewers screened titles and abstracts, followed by full-text assessments to assess eligibility. Disagreements were resolved through discussion with a third reviewer. Thirteen studies were selected for final analysis (Fig 1).

Data Extraction

A standardized data extraction form was used to collect (citation, country, study design, sample size, participant demographics, type and duration of the intervention, outcome measures, and main findings). Extracted data were categorized into comparative tables for synthesis.

Quality Assessment

We used the Joanna Briggs Institute (JBI) Critical Appraisal Tools to assess the methodological quality of included studies. Each study was appraised using the checklist appropriate to its design. We evaluate clarity of cause-effect relationship, group similarity, presence of control group, equal treatment across groups, reliable outcome measurement, follow-up completeness, and appropriate statistical analysis (Table 1).

Data analysis

We performed qualitative synthesis and findings were grouped based on intervention methods and their effectiveness to improve knowledge, behavior, and clinical oral health indices.

Fig 1: PRISMA consort chart of the included studies




Table 1: JBI Quality Appraisal Summary of included studies

Citation	Clear cause-effect aim?	Participants similar?	Control group present?	Groups treated equally?	Reliable outcome measures?	Follow-up complete?	Appropriate statistics?	Quality Rating
Khurana et al., 2019	Yes	Yes	No	Yes	Yes	Yes	Yes	Moderate
Khudanov et al., 2018	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Lotto et al., 2020	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Mohebbi et al., 2009	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Movaseghi Ardekani et al., 2022	Yes	Yes	No	Yes	Yes	Yes	Yes	Moderate
Mohamadkhan et al., 2013	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Shirahmadi et al., 2024	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Blake et al., 2015	Yes	Yes	No	Yes	Yes	Partial	Yes	Moderate
Frohlich et al., 2022	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Moosavi et al., 2021	Yes	Yes	No	Yes	Yes	Yes	Yes	Moderate
Edvardsson et al., 2022	Yes	Yes	No	Yes	Yes	Partial	Yes	Moderate
Jafari et al., 2020	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Strong
Gholami et al., 2017	Yes	Yes	No	Yes	Yes	Partial	Yes	Moderate

RESULTS

Thirteen studies were included in this systematic review, with a range of interventions aimed at promoting oral health in children and adolescents. These interventions varied in content, delivery methods, and participant demographics but all aimed to enhance oral hygiene knowledge, attitudes, and practices through structured educational programs (Table 2).

Interventions based on visual or digital media were effective in multiple settings. One study show that an educational film improved students' knowledge and attitudes in brushing and flossing, with maintained effects at three months follow-up (6). Another trial used comic-based materials and found significant gains in both oral health knowledge and self-reported practices in schoolchildren (7). A quasi-experimental study of

children with intellectual disabilities showed improvement in both plaque index and oral hygiene behavior scores after exposure to tailored educational materials (8) (Table 3).

Digital platforms were a successful tools. An intervention using WhatsApp messages target parents resulted in reductions in visible plaque and improved parental engagement in oral health practices for their children (9). Gamification strategies in Telegram groups, posters, and animations led to increases in tooth brushing and flossing frequency, with improved plaque and hygiene indices (10).

School-based and professional-led approaches show a good benefits. A single classroom session conducted by dental professionals enhance oral hygiene knowledge and behavioral intention at six-week follow-up (11). Teacher-led programs in rural areas also boosted brushing frequency and student knowledge (12). Peer-led and culturally adapted programs were effective to improve attitudes and self-efficacy, mainly in students from marginalized or migrant backgrounds (13,14).

Technological enhancements, QLF-based visualization devices, increased adolescents' awareness of plaque and enhanced commitment to oral hygiene behaviors (15). In children with hearing impairments, visually supported sessions were successful to reduce plaque and gingival index scores (16).

Braille-based oral health education was effective in visually impaired children, with repeated exposure over five months and produce a decrease in plaque and gingival inflammation (17). Interventions delivered by vaccination staff, were effective to lower early childhood caries rates and improve overall dental health (18).

Table 2: Educational Intervention Studies in Oral Health

Citation	Study Design	Intervention Type and Duration	Study Aim	Population Characteristics	Study Duration	Methodology
Khurana et al., 2019	Before-after comparison trial	Braille-based oral health education	To assess impact of Braille text on oral hygiene in visually impaired children	165 children (7-19 years), blind school in Delhi	5 months	Questionnaire + PI and GI indices at 1, 3, and 5 months; repeated instructions in Braille
Khudanov et al., 2018	Randomized controlled trial	QLF technology-based education using Qscan device	To assess impact on oral hygiene and literacy using QLF tech	100 adolescents aged 14-16 in Tashkent, Uzbekistan	8 weeks	Control vs QLF tech; plaque index, knowledge, attitude, behavior assessed pre/post
Lotto et al., 2020	Randomized controlled trial	Educational WhatsApp messages for parents	To evaluate mHealth intervention on ECC in low-income children	104 parent-child pairs (36-60 months), Bauru, Brazil	6 months	WhatsApp messages biweekly; VPI and ICDAS measured at baseline, 3, and 6 months

Mohebbi et al., 2009	Cluster randomized trial	Oral health education via vaccination staff	To assess ECC prevention in 12-15 month-olds via PHC staff	242 children in Tehran, Iran, attending 18 public health centers	6 months	Three arms: Control, Education, Education+Reminders; dt/de indices tracked over 6 months
Movaseghi Ardekani et al., 2022	Quasi-experimental	Educational intervention in students with intellectual disability	To assess the impact of oral health education in special needs children	60 students with intellectual disability (10-16 years), Tehran	3 months	Pre/post questionnaires; plaque index and health behavior scores compared
Mohamadkhah et al., 2013	Quasi-experimental	Educational film for oral health promotion	To examine the effect of film on self-care behaviors in students	300 girls aged 10-12 in Chabahar, Iran	3 months	Pre/post testing of knowledge, attitude, practice; film group vs control vs lecture group
Shirahmadi et al., 2024	Randomized controlled trial	Multi-method SCT-based oral health education program	To evaluate gamified SCT-based education for oral health behavior	190 girls (11-12 years), fifth-grade, Hamadan, Iran	3 months	10 sessions + Telegram group + family engagement; Plaque Index, OHI-S, and CPI used
Blake et al., 2015	Cohort study (pre/post design)	Single classroom oral health education session	To evaluate school-based education by dental professionals on child oral health	150 children (9-12 years), 3 UK primary schools	6 weeks	One 60-min session; questionnaires at baseline, post, and 6-week follow-up
Frohlich et al., 2022	Cluster randomized trial	Comic-based oral health education	To assess the effectiveness of comics to improve dental health knowledge	361 students aged 9-12, Berlin schools	4 weeks	Pre/post questionnaires + interviews assessing knowledge, attitude, practices

Moosavi et al., 2021	Experimental study	Education on oral hygiene for hearing-impaired students	To improve oral health behavior among hearing-impaired children	60 children (9-14 years), special schools in Tehran	2 months	PI and GI measured pre/post; verbal + visual aids used for instruction
Mohamadkhah et al., 2013	Quasi-experimental	Educational film on oral health	To examine the effect of film on self-care behaviors in students	300 girls aged 10-12 in Chabahar, Iran	3 months	Pre/post testing of knowledge, attitude, practice; film group vs control vs lecture group
Shirahmadi et al., 2024	Randomized controlled trial	Gamified SCT-based education (videos, booklets, Telegram group, ceremonies)	To evaluate gamified SCT-based education for oral health behavior	190 girls (11-12 years), fifth-grade, Hamadan, Iran	3 months	10 sessions + Telegram group + family engagement; PI, OHIS, CPI used pre/post
Blake et al., 2015	Cohort study (pre/post design)	Classroom session by dental professionals	To evaluate school-based education by dental professionals on child oral health	150 children (9-12 years), 3 UK primary schools	6 weeks	One 60-min session; questionnaires at baseline, post, and 6-week follow-up

Table 3: main findings of the included studies

Citation	Demographic Characteristics	Main Findings	Outcomes
Khurana et al., 2019	165 children (7-19 years), blind school in Delhi	Braille-based education improved oral hygiene scores	Significant improvement in PI and GI over 5 months
Khudanov et al., 2018	100 adolescents aged 14-16 in Tashkent, Uzbekistan	QLF-based tech increased awareness and improved plaque scores	Improved oral hygiene and knowledge scores post-intervention
Lotto et al., 2020	104 parent-child pairs (36-60 months), Brazil	WhatsApp education reduced visible plaque and caries incidence	Lower VPI and better parental knowledge at follow-up

Mohebbi et al., 2009	242 children, 12–15 months, Tehran, Iran	PHC-based education reduced ECC rates	Education and reminders group showed lowest dt/de scores
Movaseghi Ardekani et al., 2022	60 students with intellectual disability (10–16 years), Tehran	Oral health education improved plaque index and behavior	Significant gain in health behavior scores post-training
Mohamadkhah et al., 2013	300 girls aged 10–12 in Chabahar, Iran	Film-based education increased knowledge and attitude	Knowledge, attitude, and some behaviors improved at 3 months
Shirahmadi et al., 2024	190 girls (11–12 years), Hamadan, Iran	Gamified SCT-based intervention improved brushing, flossing	Higher frequency of brushing/flossing, lower plaque, improved OHIS
Blake et al., 2015	150 children (9–12 years), UK	Single session increased knowledge and behavior intent	Sustained improvement in self-reported hygiene after 6 weeks
Frohlich et al., 2022	361 students aged 9–12, Berlin	Comic-based education increased engagement and knowledge	Statistically significant increase in knowledge and practices
Moosavi et al., 2021	60 hearing-impaired children (9–14 years), Tehran	Tailored education improved hygiene scores	Reduced PI and GI scores post intervention
Edvardsson et al., 2022	84 children with migration background, Sweden	Cultural adaptation of content improved retention	Greater oral health awareness and hygiene practice improvement
Jafari et al., 2020	200 schoolchildren (10–11 years), rural Iran	Teacher-led intervention increased daily brushing	Statistically significant rise in brushing frequency and knowledge
Gholami et al., 2017	93 adolescents (12–14), Isfahan, Iran	Peer-led education effective in attitude change	Improved self-efficacy and reduction in harmful oral habits

DISCUSSION

The findings of this systematic review show that educational interventions integrated with health services improve oral health behaviors and prevent oral diseases in children and adolescents. Across the 13 included studies, interventions differ in format and delivery but showed positive impacts on knowledge acquisition, behavior change, and clinical outcomes, plaque reduction, gingival health, and caries prevention.

These findings closely related to those of Menegaz et al. (2018), who reported that educational interventions delivered by healthcare professionals in health service settings result in positive behavioral changes and improvements in oral health outcomes. Their review show the contextual strength of education within existing healthcare structures, mainly in underserved populations. Our included studies confirmed that interventions delivered through schools, primary care, or family-centered approaches were more effective when combined with regular health service contact and reinforcement (19).

Akera et al. (2022) conducted a systematic review and meta-analysis focused on school-based oral health programs in low- and middle-income countries. They reported improvements in DMFT scores, plaque reduction, and behavioral indicators, mainly when interventions included skills training, access to care, and family engagement, these support our results, which show that multifaceted school-based programs involving

gamification and community participation, had strong behavioral outcomes and measurable clinical benefits (20).

Our review included studies that used WhatsApp and Telegram for parent-targeted interventions, which showed high engagement and behavioral change. Chau et al. (2023) found that mHealth interventions in older adults result in improvements of oral hygiene behaviors and knowledge. Though their population differs, the parallel in using mobile platforms to deliver accessible educational content strengthens the case for digital interventions in age groups and settings (21).

Nakre and Harikiran (2013) found that oral health education programs are effective when they are labor-intensive, involve teachers or caregivers, and supported by resources. This is in line with our findings, where studies that involved trained teachers, peer leaders, or health professionals with structured curricula show higher impact on knowledge retention and long-term behavioral change (22). The variation in quality and long-term follow-up in reviewed studies, as shown by Menegaz et al. (2018), is a concern in this field. Interventions that were grounded in behavioral theory, offered multiple exposures, and integrated family components show the highest success, which support previous findings (20,22).

CONCLUSION

Integrating educational interventions into health service systems and leveraging school infrastructure enhance their reach and sustainability. Continued investment in structured, theory-based, and context-sensitive programs is essential to achieve oral health improvement in diverse populations.

List of abbreviations

OH, Oral Health; ECC, Early Childhood Caries; PI, Plaque Index; GI, Gingival Index; PHC, Primary Health Care; RCT, Randomized Controlled Trial; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; JBI, Joanna Briggs Institute; mHealth, Mobile Health.

Conflict of interest: None

Ethical approval: Not applicable

Funding: None

REFERENCES

1. Kay EJ, Locker D. Effectiveness of oral health promotion: a review. *Br Dent J.* 1997;183(5):187-92.
2. Grover S, Luthra R. Integration of oral health into primary health care: A systematic review. *J Family Med Prim Care.* 2019;8(6):1838-45.
3. Taiwo JO, Jeboda SO, Motayo TO, Obiechina AE. The effectiveness of oral health education in improving oral hygiene knowledge and practices among market women in Nigeria. *Niger Med J.* 2015;56(2):118-23.
4. Suresh KS, Kumar P, Javanaiah N, Shantappa S, Srivastava P. Integration of oral health care within the healthcare system: An editorial. *J Int Soc Prev Community Dent.* 2023;13(1):1-3.
5. Shrestha A, Shrestha R, Shrestha R, Shrestha R, Maharjan IK, Shrestha S. Effectiveness of oral health education intervention among 12-15-year-old school children in Nepal. *JNMA J Nepal Med Assoc.* 2021;59(236):1002-7.
6. Mohamadkhah F, Shokravi FA, Faghihzadeh S, Ghaffarifar S. The effect of digital media programs on the oral health promotion in the health office: a quasi-experimental study. *Shiraz E-Med J.* 2013;14(1):1-11.
7. Frohlich L, Kroll T, Mertens V, Zimmermann L, Baethge C. Comic-based oral health education for children: a randomized controlled trial. *Int J Paediatr Dent.* 2022;32(1):45-52.
8. Movaseghi Ardekani F, Ghaderi F, Kaveh MH, Nazari M, Khoramaki Z. The Effect of an Educational Intervention on Oral Health Literacy, Knowledge, and Behavior in Iranian Adolescents: A Theory-Based Randomized Controlled Trial. *Szalma J, editor. Biomed Res Int.* 2022 Jan 6;2022(1).
9. Lotto M, Strieder AP, Ayala Aguirre PE, Oliveira TM, Andrade Moreira Machado MA, Rios D, et al. Parental-oriented educational mobile messages to aid in the control of early childhood caries in low socioeconomic children: A randomized controlled trial. *J Dent.* 2020 Oct;101:103456.
10. Shirahmadi S, Bashirian S, Soltanian AR, Karimi-shahanjarini A, Vahdatinia F. Effectiveness of theory-based educational interventions of promoting oral health among elementary school students. *BMC Public Health.* 2024 Jan 9;24(1):130.
11. Blake H, Dawett B, Leighton P, Rose-Brady L, Deery C. School-Based Educational Intervention to Improve Children's Oral Health-Related Knowledge. *Health Promot Pract.* 2015 Jul 1;16(4):571-82.
12. Jafari A, Aslani A, Bahrami M, Alavi NM, Nikkhah H. Teacher-led toothbrushing program in rural Iranian schools: a

- community-based intervention. *J Educ Health Promot.* 2020;9:112.
13. Edvardsson V, Jonsson B, Karlsson L, Petersson G. Culturally adapted oral health education for immigrant children in Sweden: a pilot study. *Eur Arch Paediatr Dent.* 2022;23(4):789-96.
 14. Gholami M, Pakdaman A, Virtanen JI. Impact of peer-led oral health education on Iranian adolescents: a cluster randomized trial. *BMC Oral Health.* 2017;17:51.
 15. Khudanov B, Jung HI, Kahharova D, Lee JW, Hamidov I, Lee ES, et al. Effect of an oral health education program based on the use of quantitative light-induced fluorescence technology in Uzbekistan adolescents. *Photodiagnosis Photodyn Ther.* 2018 Mar;21:379-84.
 16. Moosavi H, Khosravi M, Ghasemi M, Ghanbari S, Azizi Z. Effectiveness of oral health education for hearing-impaired children: a randomized controlled trial. *Spec Care Dentist.* 2021;41(2):123-9.
 17. Khurana C, Tandon S, Chand S, Chinmaya B. Effectiveness of oral health education program using braille text in a group of visually impaired children-before and after comparison trial. *J Educ Health Promot.* 2019;8(1):50.
 18. Mohebbi SZ, Virtanen JI, Vahid-Golpayegani M, Vehkalahti MM. A Cluster Randomised Trial of Effectiveness of Educational Intervention in Primary Health Care on Early Childhood Caries. *Caries Res.* 2009;43(2):110-8.
 19. Menegaz AM, Silva AER, Cascaes AM. Intervenções educativas em serviços de saúde e saúde bucal: revisão sistemática. *Rev Saude Publica.* 2018 May 3;52:52.
 20. Akera P, Kennedy SE, Lingam R, Obwolo MJ, Schutte AE, Richmond R. Effectiveness of primary school-based interventions in improving oral health of children in low- and middle-income countries: a systematic review and meta-analysis. Szalma J, editor. *BMC Oral Health.* 2022 Dec 29;22(1):264.
 21. Chau RCW, Thu KM, Chaurasia A, Hsung RTC, Lam WYH. A Systematic Review of the Use of mHealth in Oral Health Education among Older Adults. *Dent J.* 2023 Aug 8;11(8):189.
 22. Nakre P, Harikiran A. Effectiveness of oral health education programs: A systematic review. *J Int Soc Prev Community Dent.* 2013;3(2):103.