

# Effects Of Digital Therapeutics Based Dual-Task Proprioceptive Training On Postural Stability, Positioning Sensation And Cognition

JaeHo Yu<sup>1</sup>

<sup>1</sup>Department of Physical therapy, Sunmoon University, Address :70, Sunmoon-ro 221 beon-gil, Tangeong-myeon, Asan-si, Chungcheongnam-do, Republic of Korea, 31460, naresa@sunmoon.ac.kr

---

## Abstract

*This study compared augmented reality (AR)-based proprioception training to traditional therapy to determine if the two tasks together were effective in improving postural stability, proprioception, and cognition. Forty-five healthy adults in their 20s were randomized into three groups: AR-based DT, AR-based proprioceptive exercise, and therapist-supervised exercise. Paired t-test and independent t-test were used to determine the within and between group effects, and the three groups were subjected to one-way ANOVA and Bonferroni's post hoc analysis. For postural stability, stability index and postural stability improved post-intervention in all groups ( $p < .05$ ), with no differences between groups ( $p > .05$ ). Positioning sensation improved in all groups ( $p < .05$ ), with no difference between groups ( $p > .05$ ). Cognitive parameters showed significant differences in recognition and calculation in all groups after the intervention ( $p < .05$ ), and no significant differences in ordering ( $p < .05$ ). Thus, AR-based interventions have shown similar effects to therapists, improving cognitive performance on both tasks, and can be selected in some cases.*

**Keywords:** Digital therapeutics, Dual Task, Proprioceptive Exercise, Postural Stability, Positioning Sensation, Cognition.

---

## I. INTRODUCTION

The prevalence of musculoskeletal (MSK) symptoms continues to rise due to population aging, changing lifestyles, and increased screen time leading to VDT (Visual Display Terminal) syndrome. While regular visits to healthcare providers are recommended for managing MSK disorders, access issues and time constraints often result in inadequate treatment and chronicity. Although various consumer-grade wearable devices (e.g., TENS units, massage tools, heat stimulators) have been developed, most provide only superficial relief and fall short in addressing the root causes of MSK symptoms.

Recent studies report that up to 37% of individuals experience plantar fasciitis at some point in their lives (Khired et al., 2022). Repeated impact on the plantar fascia can result in cumulative stress on the calcaneus, gradual degradation of the heel pad, and onset of pain, joint cartilage degeneration, and arthritis. MSK fatigue and pain impair physical equilibrium and gait performance, disrupt neuromuscular signaling, and contribute to muscle weakness, joint deformity (Park et al., 2018), and postural imbalance. During warmer seasons, increased use of low-profile footwear exacerbates plantar fasciitis, while excessive pronation heightens plantar loading, further aggravating symptoms (Kim et al., 2014).

The arch-shaped slippers used in this study differ from conventional models that often induce excessive muscle contractions leading to discomfort and fatigue. Instead, these slippers promote natural activation of medial lower leg muscles critical for knee stability during daily activities. Furthermore, the design aims to minimize mediolateral sway in joint ROM and reduce GRF during gait.

Thus, the objective of this study was to evaluate the efficacy of arch-structured slippers in preventing lower limb MSK disorders by analyzing their effects on gait-related joint ROM, GRF, and muscle activation.

## I. INTRODUCTION

A double task (DT) is performing two tasks at the same time. It requires a high level of concentration as participants must perform two tasks simultaneously. (Macpherson, 2018). When adolescents underwent DT training to perform motor and cognitive tasks, their cognitive performance improved more than those in single-task training. Additionally, athletic performance improved after performing dual rather than single tasks. However, in controlled experiments, short-term effects for single-task training were better, contrary to improved long-term effects. Moreover, the degree of DT difficulty is an important factor in promoting learning ability; the higher the degree of difficulty, the greater the improvement in athletic and

walking ability obtained after performing cognitively related DT in a study involving post-stroke gait training (Emilio et al., 2014; Bustillo-Cassero et al., 2020; Sandhiya & Yamuna 2014).

Comparing the acute and chronic effects of simultaneously performing motor and cognitive tasks as a DT in different athletes showed that motor and cognitive task performances were impaired in the short term but improved in the long term. Athletes are accustomed to situations requiring multitasking abilities. The simultaneous activation of the cerebral cortices involved in exercise and cognitive functions reduces athletic performance in acute periods, although it is enhanced in chronic periods (Veldkamp, 2019). Augmented reality (AR)-based DT training is being developed to enhance cognitive task performance. As such, researchers are exploring the relevance of using AR technology to create cognitive training tasks that are closely linked to cognitive and physical function (Zhang et al., 2021), incorporating AR into medical technology (Sesmiarni et al., 2023; Liu et al., 2019).

Rehabilitation exercise programs vary in their therapeutic effects depending on posture. In conducting an in-home exercise program, a patient unfamiliar with the correct exercise posture might perform the exercises incorrectly, consequently experiencing a negative effect. The technology developed to eliminate these negative points is precisely AR-based exercise therapy. It corrects the position and movement of the body during exercise, provides exercise progress reports, and complements the unsupervised performance of the exercise programs (Laghari et al., 2021; Rauschnabel, 2021). Furthermore, AR-based exercise programs improve walking and postural stability in patients at significantly reduced postural stability and walking abilities (Cavalcanti et al., 2019).

The effectiveness of AR-based exercise programs is comparable to that of therapist-based programs (Laghari et al., 2021; Rauschnabel, 2021). Among the studies comparing the effects of AR-based exercise therapy and exercise therapy performed by a therapist, disease comparing the effects of remotely supervised in-home AR-based and directly conducted in-hospital therapist-based postural stability training showed that mobility, dynamic, and postural stability improved in both groups. The in-hospital-trained group gained more functional improvement, although the difference was insignificant (Lee et al., 2014). In addition, using an AR program for exercise programs targeting normal adults in their 20s has been shown to be an effective treatment method, similar to PT, resulting in improvements in postural stability, as well as flexibility (Lee et al., 2022). Therefore, since no significant differences exist between AR-based non-faced and face-to-face exercise programs, remote AR technology can be used to provide low-cost, high-quality, and more economical treatment (Gandolfi et al., 2017).

In patients with subacute stroke, performing DTs during 4 weeks of postural stability training significantly improved postural stability, and auditory and short-term memory compared with conventional postural stability training (Wollesen et al., 2017). DT improved walking ability by increasing stride length and walking speed after a 12-week walking training for the elderly (Ehsani et al., 2019). In addition, in training delivered to young adults, the single-task training group showed large early improvements, whereas the DT training group showed minimal differences. When measured later, the single-task group only had improved single-task function. In contrast, the DT group showed improved motor and cognitive performance, indicating a positive effect (Liu et al., 2017).

Subsequent increases in cognitive processing were observed as the complexity of the DTs increased. However, increased central interference in DT performance interferes with cognitive and motor performance in the short term owing to distraction (Ghai et al., 2016).

Despite these findings, no studies have added a DT to proprioceptive exercises using AR. In addition, there was no training in which the DT was applied to healthy adults. Therefore, this study aimed to demonstrate that AR-based proprioceptive exercises are more effective in improving postural stability, position, and awareness when performing DT combinations.

## II. MATERIALS AND METHODS

### 1) Participants

The study was conducted on 45 healthy adults, aged 20-30, who were students at S University (Table 1). The number of samples was calculated using 'GPOWER, 3.1.9.7', we set the effect size to 0.83 to obtain the required sample size of 45 (Faul et al., 2009). The inclusion criteria for this study were no history of surgery, trauma, or neurological disease related to musculoskeletal disease within the last 6 months; no pain limiting motor performance, no cognitive impairment in the past two years, no medication use related to muscle and mental illness no vestibular organ abnormality, and agreement to

participate in the study. All participants provided informed consent after the purpose and procedures of the experiment were explained.

Table 1: Participants characteristics

	DT group(n=15)	AR group(n=15)	PT group(n=15)
Age (years)	24.53±2.77	22.73±1.73	21.64±1.49
Height (cm)	171.06±7.44	167.86±7.73	167.26±8.91
Weight (kg)	64.93±9.42	62.73±9.69	59.8±8.91

Values indicate mean ± standard deviation, DT: dual task, AR: augmented reality, PT: physical therapy.

## 2) Experiment Procedures

Figure 1 depicts the research procedures. The height and weight of all participants were measured once before the experiment. The participants were randomly assigned to three groups: AR-based proprioceptive exercise (D-ARPE) with DT, AR-based proprioceptive exercise (ARPE), and therapist-supervised exercise groups. Used UINCARE Pro for the AR program. All participants, postural stability, leg length, and proprioception were measured thrice: before, during, and after the intervention. The proprioceptive exercise program lasted for 6 weeks and was applied 2 days a week for 25 min. The participants performed 2 sets X 10 reps during the first 3 weeks, and then increased to 3 sets X 10 reps after 3 weeks. The BOSU ball was used in the intervention that was performed with shoes on for safety.




Figure 1: Experimental procedure

## 3) Measurement Tools and Methods

Postural stability, proprioception, and cognitive function were measured twice, before and after experiment using the TETRAX (tetra-ataxiometric posturography, Israel), Y-balance test, a central sensorimotor integration (CSMI) equipment, and cognitive assessment battery (CAB). All measurements and evaluations were performed by the same researcher to reduce the error margin. Measuring Postural Stability

Postural stability was measured TETRAX (tetra-ataxiometric, Israel). The participants were barefoot, and their feet were placed on the flats (A, B, C, D). Measurements were performed in the following order: stable surface with eyes open (NO), stable surface with eyes closed (NC), unstable surface with eyes open (PO), and unstable surface with eyes closed (PC). A pillow was used to create an unstable support surface

(Figure 2). Measurements were taken twice: before and after intervention, for 32 s per item. The weight distribution index (WDI), stability index (ST) were used for evaluation. ST is a measure of stability during postural change, whereas WDI represents weight load and is expressed as a percentage (Akkaya et al., 2015; Gündüz et al., 2017).



Figure 2. TETRAX

The dynamic postural stability evaluation measured the functional reach of the lower extremities using the Y-balance test. The participant stood on the center box with one lower limb. After stretching the contralateral limb in three directions (anteriorly (AT), postero-medially (PM), and posterolaterally (PL)), the distance reached were summed, and the result was divided by three times the limb length. Multiplying this quotient by 100 resulted in the composite reach distance. Before commencing, the participants had shoes off to reduce interference with postural stability. Then, the participants performed the exercise in the three directions, with three measurements each. If participants could not touch the boxes (Figure 3) or maintain their posture, the results were invalidated and remeasured (Rauh et al., 2021).



Figure 3a-d: Y-balance

Limb length was measured with the participant positioned supine on a bed. The measurer passively the ASIS to the most prominent part of the medial malleolus in cm (Hoyle and Latour, 1991). Proprioception was measured an isokinetic measuring equipment, CSMI (Human Co, USA, 2010) as follows: The measurer trained the participant to bend the joints and move their legs, and then arbitrarily set the leg positions of the participant. We then instructed participants to memorize the initial position and asked them to create an initial target angle with eyes closed (Figure 4). To reduce the error, we took three measurements and calculated the average value (Seven et al., 2019).



Figure 4: CSMI

The cognitive function was evaluated using a neurocognitive test CoTras. The CoTras consists of six short tests (cognition of speed, attention, episodic memory, visuospatial, language, and executive function).

#### 4) Intervention Method

##### Intervention Method

Of the three groups, the D-ARPE group will be subjected. In this study, the cognitive task and the peculiar training using the BOSU ball was performed simultaneously during the DT practice. Cognitive tasks included subtraction task, word memory, and ordering tasks. In the subtraction task, after the experimenter assigned the participant a numerical value for  $n$ , the participant was asked to calculate the number  $-7$  for  $n$  while simultaneously performing the BOSU ball-based training. For example, like as serial subtraction, participants repetitively perform subtraction on an arbitrarily assigned number "100-7=93, 93-7=86...". In the task of remembering words, the experimenter audibly pronounced 10 words during the proprioceptive training and then checked whether the participant remembered the words pronounced during the proprioceptive training. In the subsequent task, the participants memorized and confirmed the words and numbers presented by the experimenter during proprioceptive training (Strobach, 20191).

##### Proprioceptive Exercise Program

The exercise program is structured as follows (Table 2). AR follows AR verbal commands and visual feedback, PT follows the therapist's commands.

#### 5) Data Analysis

All statistical analyses were using the IBM SPSS 28.0.1.1. Technical statistics were used to assess the general properties. All mutations in the population yielded mean (M) and standard deviation. A paired sample t-test was used for pre-and post-variation of the intra-population exercise programs. A one-way ANOVA was used to examine the results between groups. All statistical levels ( ) were set at  $<0.05$ . Post hoc tests used Bonferroni.

### III. RESULTS

#### 1) Postural Stability

TETRAX pre-intervention measurement showed no significant differences in postural stability among all groups ( $p > 0.05$ ). Pre- and post-intervention results for each group showed within-group differences in ST during PO and PC in all groups, and significant differences in ST during NC in the -ARPE group ( $p < 0.05$ ; Table 4). No differences in ST during NO among the groups ( $p > 0.05$ ). There was no difference when measured after arbitration ( $p > 0.05$ ).

#### 2) Dynamic Postural Stability

There was no difference in the Y-balance pre-intervention measurement for PL, PM, and ANT ( $p > 0.05$ ) among all groups. Measurements within group after intervention showed differences in PL, PM, and ANT in all three groups ( $p < 0.05$ ; Table 6). Post-intervention between-group measurements showed no significant differences in ANT, PM, or PL ( $p > 0.05$ ).



### 3) Proprioception

There was no difference in proprioception among the groups in the pre-intervention measurement results ( $p > 0.05$ ). Post-intervention within-group measures showed significant differences ( $p < 0.05$ ). There was no difference between the group measures after intervention ( $p > 0.05$ ).

## IV. DISCUSSION

Poor proprioception and lack of dynamic and postural stability lead to repetitive injuries and instability, causing many problems when performing dynamic movements. Additionally, people with postural stability problems rely on proprioception. Therefore, in the process of implementing postural stability rehabilitation training, proprioceptive training is essential (Alghadir et al., 2020). Postural stability training with an instability tool such as the BOSU ball has a positive impact on postural stability, gait, and postural stability and acts effectively as proprioceptive training (Martinez-Amat et al., 2013). Research on AR-based rehabilitation training using these tools is ongoing. Four weeks of AR rehabilitation training in older adults improved postural stability, mobility, and stability of the lower extremities and reduced the risk of falls (Im et al., 2015). This means that AR rehabilitation training can improve postural stability without the need for therapist supervision. Furthermore, 4 weeks of single- and DT postural stability training in older adults with osteoporosis showed improvements in postural stability (Konak et al., 2016). These findings suggest that DT is more effective when performed continuously. The study found no difference between the D-ARPE, ARPE, and therapist-supervised exercise groups. Previous studies have demonstrated that training with unstable tools can improve

Dynamic postural stability is an important criterion in the prediction, prevention, and rehabilitation of lower-extremity musculoskeletal injuries (Powden et al., 2019). The interventions in this study increased dynamic postural stability, YBT performance, and muscle strength. In this study reported that YBT performance was affected by lower-extremity muscle strength (Lee et al., 2015a). In this study found that muscle loss around the hip reduces pelvic stability and affects the ankle and foot positions to maintain it causing an adverse effect on lower-extremity postural stability and functional movements (Jaber et al., 2018). Furthermore, in this study reported that athletes with ankle ligament injuries showed improved dynamic postural stability, muscle strength, ROM, muscular endurance, and functional ability through proprioceptive exercises with a BOSU ball (Ahern et al., 2021). Kang et al. (2015) reported that the ANT reach was correlated with ankle dorsiflexion body extension and postural stability and that hip flexion and body flexion have a positive effect on PM and PL reach. Jelinek et al. (2019) reported that knee and ankle stability and proprioception contribute to postural stability control and affect YBT. In conclusion, the body, hip, knee, and ankle play important roles in the YBT range, with the ankle and knee stability muscles in the ANT and the knee and hip stability muscles in the PM and PL being the main factors. The study's exercise program, which consisted of exercises for these muscle groups and stability, showed strong improvements in postural stability in all directions, which is consistent with results reported in previous studies. In studies involving YBT, the effects of AR-based exercises were not significantly different from conventional treatment. An 8-week Otago exercise program improved dynamic postural stability in middle-aged adults. This indicates that AR-based rehabilitation training is not different from therapist-supervised conventional exercises (Almarzouki et al., 2020). Moreover, the results of comparing the single- and DT groups showed a further improved postural stability effect in the DT group (Hoshyari et al., 2022). In contrast, this study found no significant differences among the groups, with improved results within the groups.

Proprioception plays an important role in the control of human movement. In addition, it plays an important role in joint stability, kinematic sensitivity, and postural stability control. Further, it is a key element in joint sensation, body movement, and alignment (Han et al., 2016). Roijezon et al. (2015) reported that proprioceptive deficits may appear as a result of traumatic injury to ligaments and muscle tissue and are also associated with pain disorders that develop gradually thereafter. Furthermore, Riva et al. (2019) reported that decreased proprioception was associated with excessive visual field dependence and lack of interaction with uneven ground, increasing the risk of falls. They found that the likelihood of ankle sprain recurrence in a proprioceptive training group decreased by 35% compared with that in a control group (Rivera et al., 2017). In addition, Tsauo et al. (2008) reported that 8 weeks of proprioceptive training in knee osteoarthritis patients reduced secondary injury rates. This suggests that steady proprioceptive training can improve postural stability and movement preventing traumatic injury (Relph

and Lee, 2016). Lee et al reported that, as a result of 8 weeks of single- and DT proprioceptive training in stroke patients, the DT group showed greater improvement in proprioception (Lee et al., 2015b). This suggests that DTs are more effective than single tasks. There were no differences between the groups in this study, we showed improved results when comparing before and after intervention within groups.

Routine activities, such as crossing the street, require the management of motor and cognitive tasks while processing external information. Cognitive decline often leads to reduced physical performance and gait disturbances. This change manifests in an increased risk of falls. Rave et al. (2014) reported that intervention between a single-task motor group and DT motor-cognitive group in older adults resulted in improved simultaneous attention and reduced fall risk in the DT group. Jamie et al (2017) noted that a combined physical and cognitive intervention improved cognitive function in healthy older adults with cognitive impairment and was more effective than sequential training. In addition, as a result of performing dual and single tasks in patients with brain damage, Parkinson's disease, and Alzheimer's disease, they reported that postural stability and cognition were further improved by performing DTs (Fritz et al., 2016). This suggests that performing DTs has a better effect on cognition than performing single tasks. In this study, cognitive ability and arithmetic skills improved in the DT group that underwent motor and cognitive training.

This study has some limitations. First, it targets healthy adults in their 20s; therefore, generalizing the results to all age groups should be done with caution. Second, there were limitations in the control group because they could not participate in the daily life of the experimenters. The third is UINCARE Pro recognition ability, which is an AR program. This is the delay in the process of re-recognition due to a sudden body recognition error in performing the exercise. However, in the present study, AR-based exercise and exercise performed by a physiotherapist showed similar results, and there was no significant difference between the D-ARPE group, ARPE, and therapist-based groups. Currently, it is possible to visit many rehabilitation hospitals in the vicinity due to the development of transportation means in each area. However, with many media offerings and diverse natural disasters, patients increasingly prefer in-home training. Additionally, the AR program can receive instant feedback without a therapist, improve exercise accuracy, and is suitable for use during in-home training.

## V. Conclusions

The purpose of this study was to confirm the improvement in postural stability, position sense, and cognition due to DTs when applying DTs to AR-based arbitration. The results are as follows: First, motor programs and DTs improved postural stability, cognition, and position sense. Second, the exercise programs performed in D-ARPE, ARPE, and therapist-based exercises were effective in improving postural stability and proprioception. DTs and AR had similar effects on postural stability ability and proprioception when compared with therapist-based exercises. The DT-ARPE resulted in additional cognitive improvements. AR-based interventions have been shown to have similar effects to therapists and are effective for cognition and can be an option in some cases.

## REFERENCES

- [1] physical activity level. *International journal of sports physical therapy*, 10(6), 893-900.
- [2] Alghadir, A.H., Iqbal, Z.A., Iqbal, A., Ahmed, H., & Ramteke, S.U. (2020). Effect of chronic ankle sprain on pain, range of motion, proprioception, and balance among athletes. *International journal of environmental research and public health*, 17(15), 5318. <https://doi.org/10.3390/ijerph17155318>
- [3] Almarzouki, R., Bains, G., Lohman, E., Bradley, B., Nelson, T., Alqabbani, S., & Daher, N. (2020). Improved balance in middle-aged adults after 8 weeks of a modified version of Otago Exercise Program: A randomized controlled trial. *Plos one*, 15(7), e0235734. <https://doi.org/10.1371/journal.pone.0235734>
- [4] Bustillo-Cassero, P., Cebrian-Bou, S., Cruz-Montecinos, C., Pardo, A., & García-Massó, X. (2020). Effects of a dual-task intervention in postural control and cognitive performance in adolescents. *Journal of Motor Behavior*, 52(2), 187-195.
- [5] Cavalcanti, V.C., De Santana Ferreira, M.I., Teichrieb, V., Barioni, R.R., Correia, W.F.M., & Da Gama, A.E.F. (2019). Usability and effects of text, image and audio feedback on exercise correction during augmented reality based motor rehabilitation. *Computers & Graphics*, 85, 100-110.
- [6] Deivendran, K., Hairul Anuar, H., Taran, S., & Mohamed Saat, I. (2017). Efficacy of functional rehabilitation on dynamic balance in athletes with lateral ligament ankle injury. *Research Journal of Pharmacy and Technology*, 10(12), 6-7.
- [7] Zamrini, E., Tirambulo, C., & Toosizadeh, N. (2019). The association between cognition and dual-tasking among older adults: the effect of motor function type and cognition task difficulty. *Clinical interventions in aging*, 14, 659-669. <https://doi.org/10.2147/CIA.S198697>
- [8] Emilio, E.J.M.L., Hita-Conterras, F., Jiménez-Lara, P.M., Latorre-Román, P., & Martínez-Amat, A. (2014). The association of flexibility, balance, and lumbar strength with balance ability: risk of falls in older adults. *Journal of sports science & medicine*,

13(2), 349-357.

- [9] Faul, F., Erdfelder, E., Buchner, A., & Lang, A.G. (2009). Statistical power analyses using G\* Power 3.1: Tests for correlation and regression analyses. *Behavior research methods*, 41(4), 1149-1160.
- [10] Fritz, N.E., Cheek, F.M., & Nichols-Larsen, D.S. (2015). Motor-cognitive dual-task training in neurologic disorders: a systematic review. *Journal of neurologic physical therapy: JNPT*, 39(3), 142-153.
- [11] Gandolfi, M., Geroi, C., Dimitrova, E., Boldrini, P., Waldner, A., Bonadiman, S., & Smania, multicenter, single-blind, randomized, controlled trial. *BioMed research international*, 2017.
- [12] Ghai, S., Ghai, I., & Effenberg, A.O. (2017). Effects of dual tasks and dual-task training on postural stability: a systematic review and meta-analysis. *Clinical Interventions in Aging*, 12, 557-577. <https://doi.org/10.2147/cia.s125201>
- [13] Gündüz, O.H. (2017). What Impairs Balance in Ankylosing Spondylitis? Posture or Disease Activity? *Archives of Rheumatology*, 32(3), 221-226.
- [14] Han, J., Waddington, G., Adams, R., Anson, J., & Liu, Y. (2016). Assessing proprioception: A critical review of methods. *Journal of Sport and Health Science*, 5(1), 80-90.
- [15] Hoshyari, M., Saemi, E., & Doustan, M. (2022). Motor learning and working memory in children: The role of cognitive-motor and motor-motor dual-task training. *Psihologija*, 55(4), 397-413.
- [16] Hoyle, D.A., Latour, M., & Bohannon, R.W. (1991). Intraexaminer, Interexaminer, and Interdevice Comparability of Leg Length Measurements Obtained with Measuring Tape and Metrecom. *Journal of Orthopaedic & Sports Physical Therapy*, 14(6), 263-268.
- [17] Im, D.J., Ku, J., Kim, Y.J., Cho, S., Cho, Y.K., Lim, T., Lee, H.S., Kim, H.J., & Kang, Y.J. (2015). Utility of a Three-Dimensional Interactive Augmented Reality Program for Balance and Mobility Rehabilitation in the Elderly: A Feasibility Study. *Annals of Rehabilitation Medicine*, 39(3), 462-472.
- [18] Jaber, H., Lohman, E., Daher, N., Bains, G., Nagaraj, A., Mayekar, P., Shanbhag, M., & Alameri, M. (2018). Neuromuscular control of ankle and hip during performance of the star excursion balance test in subjects with and without chronic ankle instability. *PLOS ONE*, 13(8), e0201479. <https://doi.org/10.1371/journal.pone.0201479>
- [19] Jelinek, H.F., Khalaf, K., Poilvet, J., Khandoker, A.H., Heale, L., & Donnan, L. (2019). The Effect of Ankle Support on Lower Limb Kinematics During the Y-Balance Test Using Non-linear Dynamic Measures. *Frontiers in Physiology*, 10. <https://doi.org/10.3389/fphys.2019.00935>
- [20] Kang, M., Kim, G., Kwon, O., Weon, J., Oh, J., & An, D. (2015). Relationship Between the PM&R, 7(11), 1152-1158.
- [21] Konak, H.E., Kibar, S., & Ergin, E.S. (2016). The effect of single-task and dual-task balance exercise programs on balance performance in adults with osteoporosis: a randomized controlled preliminary trial. *Osteoporosis International*, 27(11), 3271-3278.
- [22] Laghari, A.A., Jumani, A.K., Kumar, K., & Chhajro, M.A. (2021). Systematic Analysis of Virtual Reality & Augmented Reality. *International Journal of Information Engineering & Electronic Business*, 13(1), 36-43.
- [23] Lee, C.H., Kim, Y., & Lee, B.H. (2014). Augmented reality-based postural control training improves gait function in patients with stroke: Randomized controlled trial. *Hong Kong Physiotherapy Journal*, 32(2), 51-57.
- [24] Lee, D.K., Kang, M.H., Lee, T.S., & Oh, J.S. (2015a). Relationships among the Y balance test, Berg Balance Scale, and lower limb strength in middle-aged and older females. *Brazilian Journal of Physical Therapy*, 19(3), 227-234.
- [25] Lee, H., Kim, H., Ahn, M., & You, Y. (2015). Effects of proprioception training with exercise imagery on balance ability of stroke patients. *Journal of Physical Therapy Science*, 27(1), 14.
- [26] Lee, J., Yu, J., Hong, J., Lee, D., Kim, J., & Kim, S. (2022). The Effect of Augmented Reality-Based Proprioceptive Training Program on Balance, Positioning Sensation and Flexibility in Healthy Young Adults: A Randomized Controlled Trial. *Healthcare*, 10(7), 1202. <https://doi.org/10.3390/healthcare10071202>
- [27] Liu, J.C., Ke, Y.Q., Kao, Y.C., Tsai, S.C., & Lin, Y.B. (2019). A Dual-Task Authentication Mechanism Through SNMP. *Journal of Wireless Mobile Networks, Ubiquitous Computing, and Dependable Applications (JOWUA)*, 10(4), 31-45.
- [28] Liu, Y.C., Yang, Y.R., Tsai, Y.A., & Wang, R.Y. (2017). Cognitive and motor dual task gait training improve dual task gait performance after stroke - A randomized controlled pilot trial. *Scientific Reports*, 7(1). <https://doi.org/10.1038/s41598-017-04165-y>
- [29] MacPherson, S.E. (2018). Definition: Dual-tasking and multitasking. *Cortex: A Journal Devoted to the Study of the Nervous System and Behavior*, 106, 313-314.
- [30] Martínez-Amat, A., Hita-Contreras, F., Lomas-Vega, R., Caballero-Martínez, I., Álvarez, P.J., & Martínez-López, E. (2013). Effects of 12-Week Proprioception Training Program on Postural Stability, Gait, and Balance in Older Adults. *Journal of Strength and Conditioning Research*, 27(8), 2180-2188.
- [31] Powden, C.J., Dodds, T.K., & Gabriel, E.H. (2019). The reliability of the star excursion balance test and lower quarter y-balance test in healthy adults: A systematic review. *International Journal of Sports Physical Therapy*, 14(5), 683-694.
- [32] Rauschnabel, P.A. (2021). Augmented reality is eating the real-world! The substitution of physical products by holograms. *International Journal of Information Management*, 57, 102279. <https://doi.org/10.1016/j.ijinfomgt.2020.102279>
- [33] Relph, N., & Herrington, L. (2016). The effects of knee direction, physical activity and age on knee joint position sense. *The Knee*, 23(3), 393-398.
- [34] Riva, D., Fani, M., Benedetti, M.G., Scarsini, A., Rocca, F., & Mamo, C. (2019). Effects of High-Frequency Proprioceptive Training on Single Stance Stability in Older Adults: Implications for Fall Prevention. *BioMed Research International*, 1-11. <https://doi.org/10.1155/2019/2382747>
- [35] Rivera, M.J., Winkelman, Z.K., Powden, C.J., & Games, K.E. (2017). Proprioceptive Training for the Prevention of Ankle Sprains: An Evidence-Based Review. *Journal of Athletic Training*, 52(11), 1065-1067. <https://doi.org/10.4085/1062-6050-52.11.16>
- [36] Røijezon, U., Clark, N. C., & Treleaven, J. (2015). Proprioception in musculoskeletal rehabilitation. Part 1: Basic science and principles of assessment and clinical interventions. *Manual Therapy*, 20(3), 368-377.



<https://doi.org/10.1016/j.math.2015.01.008>

- [37] Sandhiya, K., & Yamuna, L. (2014). High Performance Dual-task Printed Doublet Design Loaded with Split Resonator Structures. *International Journal of Communication and Computer Technologies (IJCCCTS)*, 2(1), 64-67.
- [38] Sesmiarni, Z., Darmawati, G., Yuspita, Y.E., Yeri, S., & Ikhsan, I. (2023). Android-Based Augmented Reality: An Alternative in Mastering Tajweed for Student Learning. *Journal of Internet Services and Information Security (JISIS)*, 13(2), 30-47.
- [39] Seven, B., Cobanoglu, G., Oskay, D., & Atalay-Guzel, N. (2019). Test Retest Reliability of Isokinetic Wrist Strength and Proprioception Measurements. *Journal of Sport Rehabilitation*, 28(7), 01-06. <https://doi.org/10.1123/jsr.2018-0341>
- [40] Strobach, T. (2019). The dual-task practice advantage: Empirical evidence and cognitive mechanisms. *Psychonomic Bulletin & Review*, 27(1), 3-14.
- [41] Tait, J.L., Duckham, R.L., Milte, C.M., Main, L.C., & Daly, R.M. (2017). Influence of Sequential vs. Simultaneous Dual-Task Exercise Training on Cognitive Function in Older Adults. *Frontiers in Aging Neuroscience*, 9. <https://doi.org/10.3389/fnagi.2017.00368>
- [42] Tsao, J.Y., Cheng, P.F., & Yang, R.S. (2008). The effects of sensorimotor training on knee proprioception and function for patients with knee osteoarthritis: a preliminary report. *Clinical Rehabilitation*, 22(5), 448-457.
- [43] Van Het Reve, E., & de Bruin, E.D. (2014). Strength-balance supplemented with computerized cognitive training to improve dual task gait and divided attention in older adults: a multicenter randomized-controlled trial. *BMC geriatrics*, 14, 1-15.
- [44] Veldkamp, R., Baert, I., Kalron, A., TaRaats, J., Goetschalckx, M., Brichetto, G., Shalmoni, N., Hellinckx, P., De Weerd, N., De Wilde, D., & Feys, P. (2019). Structured Cognitive-Motor Dual Task Training Compared to Single Mobility Training in Persons with Multiple Sclerosis, a Multicenter RCT. *Journal of Clinical Medicine*, 8(12), 2177. <https://doi.org/10.3390/jcm8122177>
- [45] Wollesen, B., Schulz, S., Seydell, L., & Delbaere, K. (2017). Does dual task training improve walking performance of older adults with concern of falling?. *BMC geriatrics*, 17, 1-9.
- [46] Zhang, B., & Robb, N. (2021). A comparison of the effects of augmented reality N-Back training and traditional two-dimensional N-Back training on working memory. *SAGE Open*, 11(2), 21582440211014507. <https://doi.org/10.1177/21582440211014507>